

Regret bounds
*and their applications to online learning,
optimization and games*

Joon Kwon

March 29, 2024

Contents

1	Convexity tools	5
1.1	Preliminaries	5
1.2	Convexity	6
1.3	Subgradients	8
1.4	Legendre–Fenchel transform	9
1.5	Bregman divergences	10
1.6	Strong convexity and smoothness	11
2	UMD theory	14
2.1	Introduction	14
2.2	Regularizers	17
2.3	UMD iterates	18
2.4	Regret bounds	20
2.5	Time-dependent regularizers	20
3	Online linear optimization	23
3.1	Introduction to regret minimization	23
3.2	Dual averaging	27
3.3	Online mirror descent	30
3.4	Finite action set	37
3.5	Multi-armed bandit	40
4	Online convex optimization	43
4.1	Introduction	43
4.2	Loss linearization	44
4.3	Follow the regularized leader	45
4.4	Strongly convex losses	49
4.5	Online linear regression	51
5	Blackwell’s approachability	55
5.1	Closed convex cones	55
5.2	Framework	56
5.3	Blackwell’s algorithm	58
5.4	Regret-based approachability algorithms	59

5.5	Approachability-based regret minimization on the simplex . .	63
5.6	Further applications	66
6	Gradient methods in optimization	70
6.1	Lipschitz convex optimization	70
6.2	Smooth convex optimization	73
6.3	Nesterov's acceleration	75
6.4	Stochastic nonsmooth convex optimization	79
7	AdaGrad	81
7.1	Definitions	81
7.2	Regret bounds	82
7.3	Application to nonsmooth convex optimization	87
7.4	Application to smooth convex optimization	88
8	Monotone operators and fixed point iterations	91
8.1	Monotone operators	91
8.2	Bounded monotone operators	94
8.3	Lipschitz continuous monotone operators	96
8.4	Co-coercive operators and fixed point iterations	99
9	Regret learning in games	104
9.1	Normal-form games	104
9.2	Two-player zero-sum games	105
9.3	Regret learning	107
9.4	Optimistic regret learning	110

Foreword

As of 2024, this document contains lecture notes from a course given in Master 2 in *Université Paris–Saclay*. These are highly incomplete and constantly updated as the lectures are given.

The course proposes a unified presentation of regret minimization for adversarial online learning problems, and its application to various problems such as Blackwell’s approachability, optimization algorithms (GD, Nesterov, SGD, AdaGrad), variational inequalities with monotone operators (Extragradient, Mirror-Prox, Dual Extrapolation), fixed-point iterations (Krasnoselskii–Mann), and games. The presentation aims at being modular, so that introduced tools and techniques could easily be used to define and analyze new algorithms.

The central notion of this presentation is the *regret*, which will be analyzed using the Legendre–Fenchel transform and Bregman divergences. An excellent recent monograph on the topic of online learning is the following:

- Francesco Orabona. A modern introduction to online learning. *arXiv:1912.13213*, 2023.

Additional notable references on the topic include:

- H. Brendan McMahan. A survey of algorithms and analysis for adaptive online learning. *The Journal of Machine Learning Research*, 18(1):3117–3166, 2017,
- Elad Hazan. Introduction to online convex optimization. *Foundations and Trends® in Optimization*, 2(3-4):157–325, 2016,
- Shai Shalev-Shwartz. Online learning and online convex optimization. *Foundations and Trends in Machine Learning*, 4(2):107–194, 2011,
- Sébastien Bubeck. *Introduction to Online Optimization: Lecture Notes*. Princeton University, 2011,
- Nicolò Cesa-Bianchi and Gábor Lugosi. *Prediction, Learning, and Games*. Cambridge University Press, 2006.

Regarding convex analysis, we refer to the following classical book:

-
- R. Tyrrell Rockafellar. *Convex Analysis*. Princeton University Press, 1970.

Acknowledgements

These notes highly benefited from the help of Anatoli Judistky, discussions with Sylvain Sorin, Gilles Stoltz, Marius Potfer, Bruno Ziliotto, Jaouad Mourtada, Côme Fiegel, and the encouragements from Liliane Bel.

Chapter 1

Convexity tools

We present the basic convexity notions and tools that will be used in the subsequent chapters. Most of them are classical and are given without proof.

1.1 Preliminaries

Let $d \geq 1$. Throughout the chapter, we consider Euclidean space \mathbb{R}^d equipped with its canonical inner product denoted $\langle \cdot, \cdot \rangle$. For a set $A \subset \mathbb{R}^d$, $\text{int } A$ and $\text{cl } A$ denote its interior and closure, respectively.

Definition 1.1.1 (Domain of a function). The *domain* of a function $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ is the set

$$\text{dom } f = \left\{ x \in \mathbb{R}^d, f(x) < +\infty \right\}.$$

f is said to be *proper* if its domain is nonempty.

Definition 1.1.2 (Dual norm). Let $\|\cdot\|$ be a norm in \mathbb{R}^d . Its *dual norm* is defined as

$$\|y\|_* = \max_{\|x\| \leq 1} \langle y, x \rangle, \quad y \in \mathbb{R}^d.$$

Remark 1.1.3. The above maximum is indeed attained because for a given $y \in \mathbb{R}^d$, function $x \mapsto \langle y, x \rangle$ is continuous on the closed unit ball, which is compact. Besides, one can check that the dual norm is indeed a norm.

Proposition 1.1.4. Let $\|\cdot\|$ be a norm in \mathbb{R}^d . Then, $\|\cdot\|_{**} = \|\cdot\|$.

Example 1.1.5 (Common dual norms). In \mathbb{R}^d , ℓ_2 is its own dual norm, ℓ_p and ℓ_q (with $p, q \geq 1$ such that $1/p + 1/q = 1$) are dual of each other, and ℓ_1 and ℓ_∞ are dual of each other. If A is a positive definite matrix, the dual norm of the associated Mahalanobis norm $x \mapsto \sqrt{\langle x, Ax \rangle}$ is the Mahalanobis norm associated with A^{-1} .

Remark 1.1.6. It follows from the definition of the dual norm that for all $x, y \in \mathbb{R}^d$, $\langle y, x \rangle \leq \|y\|_* \|x\|$, which, together with the above examples recovers Cauchy-Schwarz and Hölder's inequalities.

For a function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ and $x \in \text{int dom } f$, if f is differentiable in x (resp. twice differentiable), we denote $\nabla f(x)$ its gradient at x (resp. $\nabla^2 f(x)$ its Hessian matrix at x).

1.2 Convexity

Definition 1.2.1. A set $\mathcal{X} \subset \mathbb{R}^d$ is *convex* if for all $x, x' \in \mathcal{X}$ and $\lambda \in [0, 1]$, $\lambda x + (1 - \lambda)x' \in \mathcal{X}$.

Example 1.2.2 (Unit balls). For all norms, as an immediate consequence of the triangle inequality, the unit ball is convex.

Example 1.2.3 (Simplex). Denote Δ_d the simplex in \mathbb{R}^d :

$$\Delta_d = \left\{ x \in \mathbb{R}_+^d, \sum_{i=1}^d x_i = 1 \right\},$$

which is a closed convex set. Note that it is contained in a hyperplane and therefore has empty interior.

Proposition 1.2.4 (Euclidean projection on a closed convex set). *Let $\mathcal{X} \subset \mathbb{R}^d$ be a closed convex set and $x \in \mathbb{R}^d$. Then, the Euclidean projection of x onto \mathcal{X} exists and is unique. In other words, $x' \mapsto \|x' - x\|_2^2$ admits a unique minimizer on \mathcal{X} .*

Definition 1.2.5 (Convex functions). A function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is *convex* if for all $x, x' \in \mathbb{R}^d$ and $\lambda \in [0, 1]$,

$$f(\lambda x + (1 - \lambda)x') \leq \lambda f(x) + (1 - \lambda)f(x').$$

f is *strictly convex* if the above inequality is strict for $\lambda \in (0, 1)$.

Remark 1.2.6. The convexity of a function f is closely related to the convexity of sets, as the former can be equivalently defined as the epigraph

$$\text{epi } f = \left\{ (a, x) \in \mathbb{R} \times \mathbb{R}^d, a \geq f(x) \right\}$$

being convex.

Example 1.2.7. The following functions are convex: linear functions, quadratic functions of the form $x \mapsto \langle x, Ax \rangle$ where A is a positive semi-definite matrix, the exponential, the negative logarithm, convex combinations of convex functions, the point-wise supremum of convex functions. Let $\|\cdot\|$ be a norm in \mathbb{R}^d and $a \geq 1$. Function $x \mapsto \|x\|^a$ is convex.

Proposition 1.2.8 (Jensen's inequality). *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex function, and X a random variable with values in $\text{dom } f$ so that $\mathbb{E}[X]$ exists. Then, f is measurable, $\mathbb{E}[f(X)]$ exists in $\mathbb{R} \cup \{+\infty\}$ and*

$$f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)].$$

In particular, for $n \geq 1$, $x_1, \dots, x_n \in \mathbb{R}^n$ and $\lambda_1, \dots, \lambda_n \geq 0$ such that $\lambda_1 + \dots + \lambda_n > 0$,

$$f\left(\frac{\sum_{i=1}^n \lambda_i x_i}{\sum_{i=1}^n \lambda_i}\right) \leq \frac{\sum_{i=1}^n \lambda_i f(x_i)}{\sum_{i=1}^n \lambda_i}.$$

Proposition 1.2.9 (First and second order characterizations of convexity). *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ a function with open domain.*

(i) *If f is differentiable on its domain, f is convex if, and only if, for all $x, x' \in \text{dom } f$,*

$$f(x') \geq f(x) + \langle \nabla f(x), x' - x \rangle.$$

(ii) *If f is twice differentiable on its domain, f is convex if, and only if, for all $x \in \text{dom } f$, $\nabla^2 f(x)$ is positive semi-definite.*

Proposition 1.2.10. *Let $\mathcal{X} \subset \mathbb{R}^d$ be a nonempty convex set, and $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex function, differentiable on an open set containing \mathcal{X} . Then, $x_* \in \mathcal{X}$ is a minimizer of f on \mathcal{X} if, and only if,*

$$\forall x \in \mathcal{X}, \langle \nabla f(x_*), x - x_* \rangle \geq 0.$$

Definition 1.2.11 (Lower semicontinuity). *A function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is lower semicontinuous if for all $a \in \mathbb{R}^d$, the set $\{x \in \mathbb{R}^d, f(x) \leq a\}$ is closed.*

Example 1.2.12. *Let $\mathcal{X} \subset \mathbb{R}^d$ be a convex set. The convex indicator of \mathcal{X} is the convex function defined as*

$$I_{\mathcal{X}}(x) = \begin{cases} 0 & \text{if } x \in \mathcal{X} \\ +\infty & \text{otherwise,} \end{cases}$$

which is lower semicontinuous if, and only if \mathcal{X} is closed.

Example 1.2.13 (Negative entropy on the simplex). *The function $h_{\text{ent}} : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ defined as*

$$h_{\text{ent}}(x) = \begin{cases} \sum_{i=1}^d x_i \log x_i & \text{if } x \in \Delta_d \\ +\infty & \text{otherwise,} \end{cases}$$

with convention $0 \log 0 = 0$, is a lower semicontinuous convex function. It will also be called the entropic regularizer.

Proposition 1.2.14. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper lower semicontinuous function and $\mathcal{X}_0 \subset \mathbb{R}^d$ a compact set such that $\text{dom } f \cap \mathcal{X}_0 \neq \emptyset$. Then, f attains a minimum on \mathcal{X}_0 .*

1.3 Subgradients

Definition 1.3.1 (Subgradients). Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ and $x, y \in \mathbb{R}^d$. y is a *subgradient* of f at x if for all $x' \in \mathbb{R}^d$,

$$f(x') \geq f(x) + \langle y, x' - x \rangle.$$

The set of all subgradients of f at x is called the *subdifferential* of f at x and is denoted as $\partial f(x)$.

Example 1.3.2 (Absolute value). For $f : x \mapsto |x|$ defined on \mathbb{R} , the subdifferential is given by

$$\partial f(x) = \begin{cases} \{-1\} & \text{if } x < 0 \\ [0, 1] & \text{if } x = 0 \\ \{1\} & \text{if } x > 0. \end{cases}$$

Proposition 1.3.3. Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$. A point $x_* \in \mathbb{R}^d$ is a global minimizer of f if, and only if $0 \in \partial f(x_*)$.

Proof. x_* being a global minimizer can be written

$$\forall x \in \mathbb{R}^d, \quad f(x) \geq f(x_*) + \langle 0, x - x_* \rangle,$$

in other words $0 \in \partial f(x_*)$. □

Proposition 1.3.4. Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a convex function and $x \in \text{int dom } f$. f is differentiable in x if, and only if, $\partial f(x)$ is a singleton. When this is the case, $\partial f(x) = \{\nabla f(x)\}$.

Remark 1.3.5. Even in the case of a point in the domain of a convex function, the subdifferential may be empty. Consider for instance $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ defined as:

$$f(x) = \begin{cases} -\sqrt{x} & \text{if } x \geq 0 \\ +\infty & \text{if } x < 0, \end{cases}$$

which is a proper lower semicontinuous convex function. 0 belongs to the domain of f and yet, $\partial f(0) = \emptyset$.

Proposition 1.3.6 (see e.g. Theorem 23.4 in [Roc70]). Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper convex function. If $x \notin \text{dom } f(x)$, then $\partial f(x) = \emptyset$ and if $x \in \text{int dom } f$, then $\partial f(x) \neq \emptyset$.

Proposition 1.3.7. Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper convex function, $\|\cdot\|$ a norm on \mathbb{R}^d , and $L > 0$. Then, f is L -Lipschitz in $\text{int dom } f$ with respect to $\|\cdot\|$ if, and only if:

$$\forall x \in \text{int dom } f, \quad \forall y \in \partial f(x), \quad \|y\|_* \leq L.$$

1.4 Legendre–Fenchel transform

Definition 1.4.1. Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper function. The *Legendre–Fenchel transform* (or *convex conjugate*) of f is a function $f^* : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ defined as

$$f^*(y) = \sup_{x \in \mathbb{R}^d} \{ \langle y, x \rangle - f(x) \}, \quad y \in \mathbb{R}^d.$$

Remark 1.4.2. In the above definition, for a given $y \in \mathbb{R}^d$, because f is assumed proper, quantity $\langle y, x \rangle - f(x)$ is not $-\infty$ for at least some point $x \in \mathbb{R}^d$, and therefore, the supremum is indeed a value in $\mathbb{R} \cup \{+\infty\}$.

Remark 1.4.3 (Fenchel’s inequality). It follows from the above definition that for all $x, y \in \mathbb{R}^d$, $\langle y, x \rangle \leq f(x) + f^*(y)$.

The above definition is somewhat abstract, and it may be insightful to informally examine a simple example in dimension 1 by decomposing the transformation into simpler steps. Consider $f(x) = \frac{a}{2}(x - b)^2$ for some $a, b \neq 0$, which is a differentiable convex function with finite values on \mathbb{R} . Its derivative is given by $f'(x) = a(x - b)$, which is (strictly) increasing (as would be the case as soon as f is differentiable and strictly convex), and is a bijection from its domain to its range (from \mathbb{R} to \mathbb{R} in this case). Then, the inverse $(f')^{-1}$ is also an increasing function: $(f')^{-1}(y) = y/a + b$. We then consider the following primitive function:

$$f^*(y) = \int_0^y (f')^{-1} - \min_{x \in \mathbb{R}} f(x), \quad y \in \mathbb{R},$$

which can be proved to be an alternative definition of the Legendre–Fenchel transform in this special case (although the proof is somewhat involved). As the primitive of an increasing function, f^* is also convex. Because of the inverse relation between the derivatives, this transformation $f \mapsto f^*$ is involutorial up a constant: $f^{**} = f + a$, ($a \in \mathbb{R}$). Then, it can be proved that $a = 0$, which can also be noticed graphically. An intuition that appears in this example is that the more f is curved, the less f^* is so, and vice-versa. The derivatives being inverses of each other can be interpreted as follows: f has slope y at point x if, and only if, f^* has slope x at point y . In higher dimension, this builds on the usual duality between points and hyperplanes. The Legendre–Fenchel transform indeed generalizes the above to higher dimensions, and for nondifferentiable functions. The neatest properties of e.g. Propositions 1.4.5 and 1.4.6 below are obtained for the class of proper lower semicontinuous convex functions. The derivative, which is a *function*, is then replaced by the subdifferential, which is a *correspondence*.

Proposition 1.4.4. *If $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is a proper function, f^* is lower semicontinuous and convex.*

Theorem 1.4.5 (Fenchel–Moreau). *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper function. Then, f is lower semicontinuous and convex if, and only if $f = f^{**}$. In this case, f^* is proper.*

Proposition 1.4.6. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper lower semicontinuous convex function and $x, y \in \mathbb{R}^d$. The following statements are equivalent:*

- (i) $x \in \partial f^*(y)$,
- (ii) $y \in \partial f(x)$,
- (iii) $\langle y, x \rangle = f(x) + f^*(y)$,
- (iv) $x \in \text{Arg max}_{x' \in \mathbb{R}^d} \{\langle y, x' \rangle - f(x')\}$,
- (v) $y \in \text{Arg max}_{y' \in \mathbb{R}^d} \{\langle y', x \rangle - f^*(y')\}$.

Example 1.4.7 (Norms and squared norms). Let $\|\cdot\|$ be a norm in \mathbb{R}^d and denote B its closed unit ball. Then, I_B is then a proper lower semicontinuous convex function and $I_B^* = \|\cdot\|_*$. Therefore, the involutonal property of the Legendre-Fenchel transform is an extension of the involutonal property for dual norms. Besides, if $f : x \mapsto \frac{1}{2} \|x\|^2$, then for $y \in \mathbb{R}^d$, $f^*(y) = \frac{1}{2} \|y\|_*^2$.

1.5 Bregman divergences

We now define a large class of similarity measures in \mathbb{R}^d called Bregman divergences, which in general are not distances because they may fail to be symmetric. They contain the squared Euclidean norm and the Kullback–Leibler divergence as special cases. Bregman divergences are used as an alternative geometry to the Euclidean one when it comes to e.g. defining and analyzing iterative algorithms. We present the classical definition which involves a gradient.

Definition 1.5.1. Let $\mathcal{X} \subset \mathbb{R}^d$, $f : \mathcal{X} \rightarrow \mathbb{R}$, $x \in \text{int } \mathcal{X}$ and $x' \in \mathcal{X}$ such that f is differentiable in x . Then, the *Bregman divergence* from x to x' is defined as

$$D_f(x', x) = f(x') - f(x) - \langle \nabla f(x), x' - x \rangle.$$

Remark 1.5.2. $D_f(x, x')$ is the remainder of the first order Taylor’s expansion from x to x' , and is a measure of the curvature of f between those two points. The Bregman divergence is nonnegative as soon as f is convex. In the case of a linear function f , the Bregman divergence is zero, which corresponds to the linear function having no curvature.

The above definition which requires the differentiability at starting point x is the most common. For our purposes however, we also consider the following generalization (proposed in [JKM23]) which involves a subgradient instead of a gradient.

Definition 1.5.3 (Generalized Bregman divergences). Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a function, $x, x', y \in \mathbb{R}^d$ such that $x \in \text{dom } f$ and $y \in \partial f(x)$. The *Bregman divergence* from x to x' with subgradient y is then defined as

$$D_f(x', x; y) = f(x') - f(x) - \langle y, x' - x \rangle.$$

Remark 1.5.4. The above generalized Bregman divergence may not exist even when x belongs to the domain of f , as the subdifferential may be empty. When it exists, because $x \in \text{dom } f$, it belongs to $\mathbb{R} \cup \{+\infty\}$ and is nonnegative when f is convex. When f is convex and differentiable at point x , the only subgradient at x is $\nabla f(x)$ according to Proposition 1.3.4, and the two previous definitions coincide.

Proposition 1.5.5. Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper lower semicontinuous convex function and $x, x', y, y' \in \mathbb{R}^d$ such that $y \in \partial f(x)$ and $y' \in \partial f(x')$. Then,

$$D_f(x', x; y) = D_{f^*}(y, y'; x').$$

Proof.

$$\begin{aligned} D_f(x', x; y) - D_{f^*}(y, y'; x') &= f(x') - f(x) - \langle y, x' - x \rangle \\ &\quad - f^*(y) + f^*(y') + \langle x', y - y' \rangle \\ &= \langle x', y' \rangle - \langle x, y \rangle - \langle y, x' - x \rangle + \langle x', y - y' \rangle \\ &= 0, \end{aligned}$$

where for the second equality, we applied Fenchel's identity from property (iii) in Proposition 1.4.6. \square

Example 1.5.6 (Squared Euclidean norm). Consider the squared Euclidean norm $H_2 : x \mapsto \frac{1}{2} \|x\|_2^2$, which is differentiable in \mathbb{R}^d . Then for all $x, x' \in \mathbb{R}^d$, $D_{H_2}(x', x) = \frac{1}{2} \|x' - x\|_2^2$.

Example 1.5.7 (Squared Mahalanobis norm). Let A be a symmetric positive definite matrix of size d . Consider the squared Mahalanobis norm $H_A : x \mapsto \frac{1}{2} \|x\|_A^2$, which is differentiable in \mathbb{R}^d . Then for all $x, x' \in \mathbb{R}^d$, $D_{H_A}(x', x) = \frac{1}{2} \|x' - x\|_A^2$.

1.6 Strong convexity and smoothness

We now introduce strongly convexity and smoothness which intuitively correspond to the curvature of a function being respectively bounded from below (by a positive number), and bounded from above in absolute value.

Definition 1.6.1 (Strong convexity). Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$, $\|\cdot\|$ a norm in \mathbb{R}^d and $K > 0$. f is *K -strongly convex* with respect to $\|\cdot\|$ if for all $x, x' \in \mathbb{R}^d$ and $\lambda \in [0, 1]$,

$$f(\lambda x + (1 - \lambda)x') \leq \lambda f(x) + (1 - \lambda)f(x') - \frac{K\lambda(1 - \lambda)}{2} \|x' - x\|^2.$$

Definition 1.6.2 (Smoothness). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a differentiable function, $\|\cdot\|$ a norm in \mathbb{R}^d and $L > 0$. f is L -smooth with respect to $\|\cdot\|$ if for all $x, x' \in \mathbb{R}^d$, $|D_f(x', x)| \leq \frac{L}{2} \|x' - x\|^2$, in other words,

$$|f(x') - f(x) - \langle \nabla f(x), x' - x \rangle| \leq \frac{L}{2} \|x' - x\|^2.$$

Remark 1.6.3. If f is convex, the above definition reduces to $D_f(x', x) \leq \frac{L}{2} \|x' - x\|_2^2$ ($x, x' \in \mathbb{R}^d$).

Proposition 1.6.4 (Duality between strong convexity and smoothness). Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper, lower semicontinuous convex function, $\|\cdot\|$ a norm in \mathbb{R}^d and $K > 0$. The following statements are equivalent.

(i) f is K -strongly convex with respect to $\|\cdot\|$.

(ii) For all $x, x', y \in \mathbb{R}^d$ such that $y \in \partial f(x)$, $D_f(x', x; y) \geq \frac{K}{2} \|x' - x\|^2$, in other words

$$f(x') \geq f(x) + \langle y, x' - x \rangle + \frac{K}{2} \|x' - x\|^2.$$

(iii) For all $x, x', y, y' \in \mathbb{R}^d$ such that $y \in \partial f(x)$ and $y' \in \partial f(x')$,

$$\langle y' - y, x' - x \rangle \geq K \|x' - x\|^2.$$

(iv) For all $x, x', y, y' \in \mathbb{R}^d$ such that $y \in \partial f(x)$ and $y' \in \partial f(x')$,

$$\langle y' - y, x' - x \rangle \leq \frac{1}{K} \|y' - y\|_*^2.$$

(v) For all $x, x', y, y' \in \mathbb{R}^d$ such that $y \in \partial f(x)$ and $y' \in \partial f(x')$, $D_f(x', x; y) \leq \frac{1}{2K} \|y' - y\|_*^2$, in other words

$$f(x') \leq f(x) + \langle y, x' - x \rangle + \frac{1}{2K} \|y' - y\|_*^2.$$

(vi) f^* is differentiable on \mathbb{R}^d and $1/K$ -smooth with respect to $\|\cdot\|_*$.

Corollary 1.6.5. Let $K > 0$, $\|\cdot\|$ a norm on \mathbb{R}^d and $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ a proper lower semicontinuous function which we assume K -strongly convex with respect to $\|\cdot\|$. Then, for all $y, y' \in \mathbb{R}^d$,

$$D_{h^*}(y', y) \leq \frac{1}{2K} \|y' - y\|_*^2.$$

Proposition 1.6.6 (Second order characterization of strong convexity). Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a twice differentiable function, $\|\cdot\|$ a norm in \mathbb{R}^d and $K > 0$. Then, f is K -strongly convex with respect to $\|\cdot\|$ if, and only if,

$$\forall x \in \mathbb{R}^d, \forall u \in \mathbb{R}^d, \quad \langle u, \nabla^2 f(x)u \rangle \geq K \|u\|^2.$$

Proposition 1.6.7 (First and second order characterizations of smoothness). *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a differentiable function, $\|\cdot\|$ a norm in \mathbb{R}^d and $L > 0$.*

- (i) *f is L -smooth if, and only if, ∇f is L -Lipschitz with respect to $\|\cdot\|$.*
- (ii) *Moreover, if f is twice differentiable, f is L -smooth with respect to $\|\cdot\|$ if, and only if,*

$$\forall x \in \mathbb{R}^d, \forall u \in \mathbb{R}^d, \quad |\langle u, \nabla^2 f(x)u \rangle| \leq L \|u\|^2.$$

Corollary 1.6.8. *The squared Euclidean norm $h_2 : x \mapsto \frac{1}{2} \|x\|_2^2$ is 1-strongly convex and 1-smooth with respect to $\|\cdot\|_2$.*

Corollary 1.6.9. *Let A be a symmetric positive definite matrix of size d . The associated squared Mahalanobis norm $x \mapsto \frac{1}{2} \|x\|_A^2$ is 1-strongly convex and 1-smooth for $\|\cdot\|_A$.*

Proposition 1.6.10. *For $p \in (1, 2)$, the squared ℓ_p norm $h_p : x \mapsto \frac{1}{2} \|x\|_p^2$ is $(p-1)$ -strongly convex with respect to ℓ_p .*

Proposition 1.6.11. *The negative entropy h_{ent} is 1-strongly convex with respect to ℓ_1 .*

Proposition 1.6.12. *A proper lower semicontinuous strongly convex function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ admits a unique minimizer on \mathbb{R}^d .*

Chapter 2

UMD theory

Throughout the chapter, \mathcal{X} is a nonempty closed convex set of \mathbb{R}^d .

2.1 Introduction

The goal of this chapter is to introduce a general scheme for defining a sequence of iterates $(x_t)_{t \geq 0}$ in \mathcal{X} based on another sequence $(u_t)_{t \geq 0}$ in \mathbb{R}^d , where for each $t \geq 0$, vector u_t is used in the update from x_t to x_{t+1} . General properties are then established. All algorithms and guarantees from the following chapters will be derived using this general approach, called UMD for *unified mirror descent*. To get some taste and intuition, we first examine a few simple special cases before presenting our general theory.

The simplest update is given by

$$x_{t+1} = x_t + u_t, \quad t \geq 0,$$

and is already of great interest, as it contains as special cases gradient descent (where $u_t = -\gamma_t \nabla f(x_t)$ is then a step in the opposite direction of the gradient of some objective function), as well as its stochastic counterpart (SGD). Such a sequence satisfies the following elementary result.

Proposition 2.1.1. *For all $t \geq 0$ and $x \in \mathbb{R}^d$,*

$$\langle u_t, x - x_t \rangle = \frac{1}{2} \|x - x_t\|_2^2 - \frac{1}{2} \|x - x_{t+1}\|_2^2 + \frac{1}{2} \|u_t\|_2^2.$$

Consequently, for all $T \geq 0$,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle = \frac{1}{2} \|x - x_0\|_2^2 - \frac{1}{2} \|x - x_{T+1}\|_2^2 + \frac{1}{2} \sum_{t=0}^T \|u_t\|_2^2.$$

Proof. Let $t \geq 1$. Using the definition of x_{t+1} ,

$$\|x_{t+1} - x\|_2^2 = \|x_t + u_t - x\|_2^2 = \|x_t - x\|_2^2 + 2 \langle u_t, x_t - x \rangle + \|u_t\|_2^2,$$

and the result follows. \square

For instance, the classical convergence guarantees about (stochastic) gradient descent in various settings, are consequences of the above identity. The quantity $\sum_{t=0}^T \langle u_t, x - x_t \rangle$ is called the *regret*, but the corresponding interpretation will be presented in the next chapter only. Some intuition about the above can be obtained through a continuous-time counterpart:

$$\text{if } \frac{d\tilde{x}_t}{dt} = \tilde{u}_t, \quad \text{then } \frac{d}{dt} \left(\frac{1}{2} \|x - \tilde{x}_t\|_2^2 \right) = \langle \tilde{u}_t, \tilde{x}_t - x \rangle.$$

Therefore, going back to discrete-time, one can interpret the difference

$$\frac{1}{2} \|x - x_t\|_2^2 - \frac{1}{2} \|x - x_{t+1}\|_2^2$$

as a discrete-time derivative, and the term $\frac{1}{2} \|u_t\|_2^2$ —which does not appear in continuous-time—as a discretization error.

The quantity $\frac{1}{2} \|x - x_t\|_2^2$ also appears in the following alternative expression which will inspire generalizations.

Proposition 2.1.2. *For $t \geq 0$, $x_{t+1} = x_t + u_t$ if, and only if,*

$$x_{t+1} = \arg \min_{x \in \mathbb{R}^d} \left\{ -\langle u_t, x \rangle + \frac{1}{2} \|x - x_t\|_2^2 \right\}.$$

The above expression is an *incremental* point of view, in the sense that the next iterate x_{t+1} is written as a function of previous iterate x_t and vector u_t only. Another equivalent formulation is the *cumulative* one:

$$x_{t+1} = x_0 + \sum_{s=0}^t u_s,$$

where the next iterate x_{t+1} is only a function of the sum of the previous vectors u_s ($1 \leq s \leq t$) (plus the initial point x_0). These two points of view will yield different extensions below.

We now turn to a constrained setting where we need the iterates to all lie in a given nonempty closed convex set $\mathcal{X} \subset \mathbb{R}^d$. Then, the definition of the iterates can be adapted by adding a projection step onto \mathcal{X} with respect to the Euclidean distance. The projected gradient descent algorithm is a special case. Then, the iterates satisfy the following *regret bound*, where the comparison point x must lie in \mathcal{X} .

Proposition 2.1.3. *For $t \geq 0$, $x_{t+1} = \arg \min_{x \in \mathcal{X}} \|x_t + u_t - x\|_2^2$ if, and only if,*

$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \left\{ -\langle u_t, x \rangle + \frac{1}{2} \|x - x_t\|_2^2 \right\}.$$

In that case, for all $x \in \mathcal{X}$,

$$\langle u_t, x - x_t \rangle \leq \frac{1}{2} \|x - x_t\|_2^2 - \frac{1}{2} \|x - x_{t+1}\|_2^2 + \frac{1}{2} \|u_t\|_2^2.$$

Another possibility for constraining the iterates in \mathcal{X} is the following, where the vector u_t is not added to the current iterate x_t as above, but to the point before the projection onto \mathcal{X} .

Proposition 2.1.4. *If $(x_t)_{t \geq 0}$ and $(y_t)_{t \geq 0}$ satisfy*

$$y_{t+1} = y_t + u_t \quad \text{and} \quad x_{t+1} = \arg \min_{x \in \mathcal{X}} \|y_{t+1} - x\|_2^2, \quad t \geq 0,$$

then for all $t \geq 0$,

$$\langle u_t, x - x_t \rangle = \frac{1}{2} \|x - x_t\|_2^2 - \frac{1}{2} \|x - x_{t+1}\|_2^2 + \frac{1}{2} \|u_t\|_2^2.$$

We now go back to the unconstrained case ($\mathcal{X} = \mathbb{R}^d$) and consider the following generalization.

Proposition 2.1.5. *For a positive definite matrix $A \in \mathbb{R}^{d \times d}$ and $t \geq 0$, $x_{t+1} = x_t + A^{-1}u_t$ if, and only if*

$$x_{t+1} = \arg \min_{x \in \mathbb{R}^d} \left\{ -\langle u_t, x \rangle + \frac{1}{2} \|x - x_t\|_A^2 \right\}.$$

In that case, for all $x \in \mathbb{R}^d$,

$$\langle u_t, x - x_t \rangle = \frac{1}{2} \|x - x_t\|_A^2 - \frac{1}{2} \|x - x_{t+1}\|_A^2 + \frac{1}{2} \|u_t\|_{A^{-1}}^2,$$

where $\|x\|_A = \sqrt{\langle x, Ax \rangle}$ and $\|y\|_{A^{-1}} = \sqrt{\langle y, A^{-1}y \rangle}$.

The last example considers the simplex $\mathcal{X} = \Delta_d$, and appears at first sight to be quite different from the above.

Proposition 2.1.6. *If for $t \geq 0$,*

$$x_{t+1} = \left(\frac{x_{t,i} \exp(u_{t,i})}{\sum_{j=1}^d x_{t,j} \exp(u_{t,j})} \right)_{1 \leq i \leq d},$$

then for all $x \in \Delta_d$,

$$\langle u_t, x - x_t \rangle = \text{KL}(x, x_t) - \text{KL}(x, x_{t+1}) + \log \left(\sum_{i=1}^d x_{t,i} \exp(u_{t,i}) - \langle u_t, x_t \rangle \right),$$

where $\text{KL}(x', x) = \sum_{i=1}^d x'_i \log(x'_i/x_i)$ denotes the Kullback–Leibler divergence.

2.2 Regularizers

Definition 2.2.1. A function $h : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is an *pre-regularizer* on \mathcal{X} if it is strictly convex, lower semicontinuous, and if $\text{cl dom } h = \mathcal{X}$. Moreover, if $\text{dom } h^* = \mathbb{R}^d$, then h is said to be an *regularizer* on \mathcal{X} .

Remark 2.2.2. A regularizer is proper (because \mathcal{X} is nonempty), convex, and lower semicontinuous. In particular, Proposition 1.4.6 applies.

The following proposition gives several sufficient conditions for the condition $\text{dom } h^* = \mathbb{R}^d$ to be satisfied.

Proposition 2.2.3. *Let h be an pre-regularizer on \mathcal{X} .*

- (i) *If \mathcal{X} is compact, then h is a regularizer on \mathcal{X} .*
- (ii) *If h is differentiable on $\mathcal{D}_h := \text{int dom } h$ and $\nabla h(\mathcal{D}_h) = \mathbb{R}^d$, then h is a regularizer on \mathcal{X} .*
- (iii) *If h is strongly convex, then h is a regularizer on \mathcal{X} .*

Proof. Let $y \in \mathbb{R}^d$. For each of the three assumptions, let us prove that $h^*(y)$ is finite. This will prove that $\text{dom } h^* = \mathbb{R}^d$.

- (i) Because $\text{cl dom } h = \mathcal{X}$ by definition of a pre-regularizer, we have:

$$h^*(y) = \sup_{x \in \mathbb{R}^d} \{\langle y, x \rangle - h(x)\} = \sup_{x \in \mathcal{X}} \{\langle y, x \rangle - h(x)\}.$$

Besides, the function $x \mapsto \langle y, x \rangle - h(x)$ is upper semicontinuous and therefore, according to Proposition 1.2.14, attains a maximum on \mathcal{X} because \mathcal{X} is assumed to be compact. Therefore $h^*(y) < +\infty$.

- (ii) Because $\nabla h(\mathcal{D}_h) = \mathbb{R}^d$ by assumption, there exists $x \in \mathcal{D}_h$ such that $\nabla h(x) = y$. Then, by Proposition 1.4.6, $h^*(y) = \langle y, x \rangle - h(x) < +\infty$.
- (iii) The function $x \mapsto \langle y, x \rangle - h(x)$ is the opposite of a strongly convex lower semicontinuous function on \mathbb{R}^d and therefore admits a maximum by Proposition 1.6.12. Therefore, $h^*(y) < +\infty$.

□

Proposition 2.2.4 (Differentiability of h^*). *Let h be a regularizer on \mathcal{X} . Then, h^* is differentiable on \mathbb{R}^d , ∇h^* takes values in $\text{dom } h \subset \mathcal{X}$, and*

$$\nabla h^*(y) = \arg \max_{x \in \mathbb{R}^d} \{\langle y, x \rangle - h(x)\}.$$

Proof. Let $y \in \mathbb{R}^d$. Because $\text{dom } h^* = \mathbb{R}^d$, the subdifferential $\partial h^*(y)$ is nonempty by Proposition 1.3.6, $\partial h^*(y)$ is the set of maximizers of function $x \mapsto \langle y, x \rangle - h(x)$, which is strictly concave. Therefore, the maximizer belongs to $\text{dom } h$, is unique, and thus h^* is differentiable at y by Proposition 1.3.4. □

Proposition 2.2.5 (Euclidean regularizer). *The Euclidean regularizer on \mathcal{X} , defined as*

$$h_2(x) = \frac{1}{2} \|x\|_2^2 + I_{\mathcal{X}}(x), \quad x \in \mathbb{R}^d,$$

is a regularizer on \mathcal{X} and ∇h^ is the Euclidean projection onto \mathcal{X} , in other words:*

$$\nabla h_2^*(y) = \arg \min_{x \in \mathcal{X}} \|y - x\|.$$

In particular, in the unconstrained case $\mathcal{X} = \mathbb{R}^d$, $\nabla h_2^(y) = y$ for all $y \in \mathbb{R}^d$.*

Proof. □

Proposition 2.2.6 (Entropic regularizer). *The entropic regularizer*

$$h_{\text{ent}}(x) = \begin{cases} \sum_{i=1}^d x_i \log x_i & \text{if } x \in \Delta_d \\ +\infty & \text{otherwise,} \end{cases}$$

is a regularizer on Δ_d and

$$\nabla h_{\text{ent}}^*(y) = \left(\frac{\exp(y_i)}{\sum_{j=1}^d \exp(y_j)} \right)_{1 \leq i \leq d}, \quad y \in \mathbb{R}^d.$$

Proof. □

2.3 UMD iterates

Definition 2.3.1. Let h be a regularizer on \mathcal{X} and $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d . A sequence $((x_t, y_t))_{t \geq 0}$ in $\mathbb{R}^d \times \mathbb{R}^d$ is a sequence of *UMD iterates* associated with regularizer h and *dual increments* $(u_t)_{t \geq 0}$ if for all $t \geq 0$,

- (i) $y_t \in \partial h(x_t)$,
- (ii) $x_{t+1} = \nabla h^*(y_t + u_t)$.

Remark 2.3.2. By Proposition 1.4.6, property (ii) is equivalent to $y_t + u_t \in \partial h(x_{t+1})$.

Remark 2.3.3. For each $t \geq 0$, $x_t \in \text{dom } h \subset \mathcal{X}$, because otherwise $\partial h(x_t)$ would be empty by Proposition 1.3.6 and could not contain y_t .

Definition 2.3.4. Let h be a regularizer on \mathcal{X} and $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d . A sequence $((x_t, y_t))_{t \geq 0}$ in $\mathbb{R}^d \times \mathbb{R}^d$ is a sequence of *strict UMD iterates* associated with h and $(u_t)_{t \geq 0}$ if for all $t \geq 0$,

- (I) $y_t \in \partial h(x_t)$,
- (II) $\forall x \in \mathcal{X}, \langle y_t + u_t - y_{t+1} | x - x_{t+1} \rangle \leq 0$.

Proposition 2.3.5. *Let $((x_t, y_t))_{t \geq 0}$ be a sequence of strict UMD iterates defined as above. Then for all $t \geq 0$, $x_{t+1} = \nabla h^*(y_t + u_t)$ and thus $((x_t, y_t))_{t \geq 0}$ are UMD iterates.*

Proof. Let $t \geq 0$ and let us equivalently prove that $y_t + u_t \in \partial h(x_{t+1})$. Let $x \in \mathbb{R}^d$. If $x \notin \text{dom } h$,

$$+\infty = h(x) - h(x_{t+1}) \geq \langle y_t + u_t, x - x_{t+1} \rangle.$$

If $x \in \text{dom } h$, using assumption (II) and the fact that $y_{t+1} \in h(x_{t+1})$,

$$h(x) - h(x_{t+1}) \geq \langle y_{t+1}, x - x_{t+1} \rangle \geq \langle y_t + u_t, x - x_{t+1} \rangle.$$

Therefore, $y_t + u_t \in \partial h(x_{t+1})$, in other words $x_{t+1} = \nabla h^*(y_t + u_t)$. \square

Example 2.3.6 (Euclidean regularizer). Denote $\Pi_{\mathcal{X}}$ the Euclidean projection onto \mathcal{X} and consider the Euclidean regularizer on \mathcal{X} : $h = \frac{1}{2} \|\cdot\|_2^2 + I_{\mathcal{X}}$ and $x_0 \in \mathbb{R}^d$.

- If $x_{t+1} = \Pi_{\mathcal{X}}(x_t + u_t)$ for all $t \geq 0$, then $((x_t, x_t))_{t \geq 0}$ can be proved to be a sequence of strict UMD iterates.
- If $y_{t+1} = y_t + u_t$ and $x_{t+1} = \Pi_{\mathcal{X}}(y_{t+1})$ for all $t \geq 0$, then $((x_t, y_t))_{t \geq 0}$ can also be proved to be a sequence of strict UMD iterates.

Remark 2.3.7 (Non-unicity of strict UMD iterates). As already seen in the above example, an interesting character of (strict) UMD iterates is that for a given sequence $(u_t)_{t \geq 1}$ of dual increments and initial points (x_0, y_0) such that $y_0 \in \partial h(x_0)$, there may be several possible strict UMD iterates. Here is a simple and explicit example. Consider $d = 1$, $\mathcal{X} = [0, 1]$, $h(x) = \frac{1}{2}x^2 + I_{\mathcal{X}}(x)$, $(x_0, y_0) = (1, 1)$ and $u_t = (-1)^t$ for $t \geq 0$. Then, one can verify that $((1, \frac{3+(-1)^t}{2}))_{t \geq 0}$ is a strict UMD sequence, and so is $((x_t, y_t))_{t \geq 0}$ where $x_t = y_t = \frac{1+(-1)^{t+1}}{2}$ for $t \geq 1$.

Remark 2.3.8 (Existence of strict UMD iterates). As soon as regularizer h and sequence of dual increments $(u)_{t \geq 0}$ are given, we can see that associated strict UMD iterates always exist. Indeed, from the definition of a regularizer, $\text{dom } h^* = \mathbb{R}^d$ and thus one can choose any $y_0 \in \mathbb{R}^d$ and consider $x_0 := \nabla h^*(y_0)$, which satisfies $y_0 \in \partial h(x_0)$. Then, for $t \geq 0$, one can consider $y_{t+1} := y_t + u_t$ which indeed satisfies variational condition (II), and then define $x_{t+1} := \nabla h^*(y_{t+1})$, which ensures $y_{t+1} \in \partial h(x_{t+1})$, as required by (i).

Remark 2.3.9 (Alternative notation for strict UMD iterates). For a given regularizer h , let $\Pi_h : \mathbb{R}^d \rightrightarrows \mathcal{X} \times \mathbb{R}^d$ be a set-valued mapping defined as follows. For $y_0 \in \mathbb{R}^d$, $\Pi_h(y_0)$ is the set of couples (x, y) satisfying

$$x = \nabla h^*(y_0), \quad y \in \partial h(x), \quad \text{and} \quad \forall x' \in \mathcal{X}, \quad \langle y_0 - y, x' - x \rangle \leq 0.$$

Then, one can verify that $((x_t, y_t))_{t \geq 0}$ is a strict UMD sequence associated with h and given sequence $(u_t)_{t \geq 0}$ if, and only if, $y_0 \in \partial h(x_0)$ and

$$(x_{t+1}, y_{t+1}) \in \Pi_h(y_t + u_t), \quad t \geq 0.$$

2.4 Regret bounds

Lemma 2.4.1 (UMD lemma). *Let h be a regularizer on \mathcal{X} , $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , and $((x_t, y_t))_{t \geq 0}$ a sequence of UMD iterates associated with regularizer h and dual increments $(u_t)_{t \geq 0}$ and $x \in \text{dom } h$. Consider notation*

$$D_t = D_h(x, x_t; y_t), \quad D'_t = D_h(x_{t+1}, x_t; y_t), \quad D_t^* = D_{h^*}(y_t + u_t; y_t), \quad t \geq 0.$$

(i) *Then for all $t \geq 1$,*

$$\langle u_t, x - x_{t+1} \rangle = D_t - D_{t+1} - D'_t + \langle y_t + u_t - y_{t+1}, x - x_{t+1} \rangle,$$

and

$$\langle u_t, x - x_t \rangle = D_t - D_{t+1} + D_t^* + \langle y_t + u_t - y_{t+1}, x - x_{t+1} \rangle.$$

(ii) *Moreover, $((x_t, y_t))_{t \geq 0}$ are strict UMD iterates, then for all $t \geq 0$,*

$$\langle u_t, x - x_{t+1} \rangle \leq D_t - D_{t+1} - D'_t,$$

and

$$\langle u_t, x - x_t \rangle \leq D_t - D_{t+1} + D_t^*.$$

(iii) *Besides, if h is K -strongly convex with respect to some norm $\|\cdot\|$ and some $K > 0$, then for all $t \geq 0$,*

$$D_t^* \leq \frac{1}{2K} \|u_t\|_*^2.$$

Proof. The first identity from (i) can be verified by writing explicitly the difference between both sides and simplifying. The second identity follows from noticing that for all $t \geq 0$,

$$\langle u_t, x_{t+1} - x_t \rangle = D'_t + D_h(x_t, x_{t+1}; y_t + u_t) = D'_t + D_t^*,$$

where the second equality comes from Proposition 1.5.5; and adding to the first equality. Equalities in (ii) are an immediate consequence of (i). iii follows from Proposition (1.6.4). \square

2.5 Time-dependent regularizers

Definition 2.5.1. Let $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ be a sequence of regularizers and $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d . An associated sequence $((x_t, y_t))_{t \geq 0}$ in $\mathbb{R}^d \times \mathbb{R}^d$ of UMD iterates satisfy for all $t \in \mathbb{N}$,

(i) $y_t \in \partial h_t(x_t)$,

$$(ii) \quad x_{t+1} = \nabla h_{t+1/2}^*(y_t + u_t).$$

Lemma 2.5.2 (UMD lemma with time-dependent regularizers). *Let $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ be a sequence of regularizers, $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , $((x_t, y_t))_{t \geq 1}$ associated UMD iterates, and $x \in \bigcap_{t \in \frac{1}{2}\mathbb{N}} \text{dom } h_t$. For each $t \in \mathbb{N}$, consider notation*

- $D_t = D_{h_t}(x, x_t; y_t)$,
- $D'_t = h_{t+1/2}(x_{t+1}) - h_t(x_t) - \langle y_t, x_{t+1} - x_t \rangle$,
- $D_t^* = h_{t+1/2}^*(y_t + u_t) - h_t^*(y_t) - \langle u_t, x_t \rangle$,
- For $x' \in \text{dom } h_t$ (resp. $\text{dom } h_{t+1/2}$), $\Delta h_t(x') = h_{t+1/2}(x') - h_t(x')$ (resp. $\Delta h_{t+1/2}(x') = h_{t+1}(x') - h_{t+1/2}(x')$),
- $D_{t+1/2}^\Delta = \Delta h_{t+1/2}(x) - \Delta h_{t+1/2}(x_{t+1}) - \langle y_{t+1} - y_t - u_t, x - x_{t+1} \rangle$.

(i) Then for all $t \in \mathbb{N}$,

$$\langle u_t, x - x_{t+1} \rangle = D_t - D_{t+1} - D'_t + D_{t+1/2}^\Delta + \Delta h_t(x),$$

and

$$\langle u_t, x - x_t \rangle = D_t - D_{t+1} + D_t^* + D_{t+1/2}^\Delta + \Delta h_t(x).$$

(ii) If $\Delta h_t = 0$ for a given $t \in \mathbb{N}$, then

$$\langle u_t, x - x_{t+1} \rangle = D_t - D_{t+1} - D'_t + D_{t+1/2}^\Delta,$$

and

$$\langle u_t, x - x_t \rangle = D_t - D_{t+1} + D_t^* + D_{t+1/2}^\Delta.$$

(iii) If $\Delta h_{t+1/2} = 0$ and $\langle y_{t+1} - y_t - u_t, x - x_{t+1} \rangle \geq 0$ for a given $t \in \mathbb{N}$, then

$$\langle u_t, x - x_{t+1} \rangle \leq D_t - D_{t+1} - D'_t + h_{t+1}(x) - h_t(x),$$

and

$$\langle u_t, x - x_t \rangle \leq D_t - D_{t+1} + D_t^* + h_{t+1}(x) - h_t(x).$$

(iv) If for a given $t \in \mathbb{N}$, $h_{t+1/2} \geq h_t$ and h_t is K_t -strongly convex with respect to some norm $\|\cdot\|$ and some $K_t > 0$, then

$$D_t^* \leq \frac{1}{2K_t} \|u_t\|_*^2.$$

Proof. The first equality in (i) can be proved by merely simplifying. For the second inequality, we notice that

$$\begin{aligned}
\langle u_t, x_t - x_{t+1} \rangle &= D'_t + h_t(x_t) - h_{t+1/2}(x_{t+1}) - \langle y_t + u_t, x_t - x_{t+1} \rangle \\
&= D'_t + \langle y_t, x_t \rangle - h_t^*(y_t) - \langle y_t + u_t, x_{t+1} \rangle + h_{t+1/2}^*(y_t + u_t) \\
&\quad - \langle y_t + u_t, x_t - x_{t+1} \rangle \\
&= D'_t + D_t^*,
\end{aligned}$$

where for the second equality, we used Fenchel's identity from Proposition 1.4.6. Adding the above to the first equality give the second equality in (i). Then, (ii) and (iii) are easy consequences.

Let us prove (iv). The assumption $h_{t+1/2} \geq h_t$ and the definition of the Legendre–Fenchel transform immediately implies $h_{t+1/2}^* \leq h_t^*$. Therefore,

$$D_t^* = h_{t+1/2}^*(y_t + u_t) - h_t^*(y_t) - \langle u_t, x_t \rangle \leq D_{h^*}(y_t + u_t, y_t) \leq \frac{1}{2K_t} \|u_t\|_*^2.$$

□

Chapter 3

Online linear optimization

This chapter first presents the topic of regret minimization in sequential decision problems and then makes the connection with online learning and optimization through the frameworks of online linear optimization and online convex optimization. Several important families of algorithms (dual averaging, mirror descent, follow the regularized leader) are introduced and analyzed using UMD theory from Chapter 2.

Throughout the chapter, \mathcal{X} is a nonempty closed convex set, and $\Pi_{\mathcal{X}}$ denotes the Euclidean projection onto \mathcal{X} . For a function $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$, we denote $\mathcal{D}_f = \text{int dom } f$.

3.1 Introduction to regret minimization

Let us first consider a simple sequential decision problem where the Decision Maker chooses its actions in the finite set $\{1, \dots, d\}$, possibly at random. At step $t \geq 0$,

- the Decision Maker chooses $x_t \in \Delta_d$,
- Nature chooses and reveals *payoff vector* $u_t \in [0, 1]^d$,
- i_t is a random element in $\{1, \dots, d\}$ drawn according to distribution x_t and revealed to the Decision Maker,
- the Decision Maker obtains payoff u_{t,i_t} .

The choice of x_t by the Decision Maker may depend on all past information known to him, meaning $(x_0, u_0, i_0, \dots, x_{t-1}, u_{t-1}, i_{t-1})$. Similarly, the choice of u_t by Nature may depend on all past information including x_t : $(x_0, u_0, i_0, \dots, x_{t-1}, u_{t-1}, i_{t-1}, x_t)$.

In a restrictive variant of this problem, called *multi-armed bandit* and which will be considered later, the decision maker only observes the actually obtained payoff u_{t,i_t} , and not the whole payoff vector u_t .

The Decision Maker wishes to maximize its cumulative payoff $\sum_{t=0}^T u_{t,i_t}$. More specifically, we aim at constructing decision rules for the Decision Maker (which we will simply call *algorithms*) that offer some *worst-case guarantee* on the cumulative payoff which holds *for all possible sequence* $(u_t)_{t \geq 0}$ chosen by Nature. Therefore, the guarantee must be relative to the sequence of payoff vectors $(u_t)_{t \geq 0}$. One possible type of guarantee is an upper bound on the *regret*, defined as

$$\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} - \sum_{t=0}^T u_{t,i_t},$$

which compares the actual cumulative payoff with the best cumulative payoff that would have been obtained¹ by constantly choosing a given element $i \in \{1, \dots, d\}$, meaning $i_t = i$ for all $t \geq 0$.

For a given sequence of payoff vectors $(u_t)_{t \geq 0}$, the above regret is a random variable (because for each $t \geq 0$, i_t is a random variable), and we are interested in analyzing its expectation. Using the law of total expectation, we can write

$$\begin{aligned} \mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} - \sum_{t=0}^T u_{t,i_t} \right] &= \mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} - \sum_{t=0}^T \mathbb{E}[u_{t,i_t} | x_t] \right] \\ &= \mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} - \sum_{t=0}^T \langle u_t, x_t \rangle \right]. \end{aligned} \quad (3.1)$$

Besides, because the maximum of linear function on a convex compact set is attained at its edges,

$$\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} = \max_{1 \leq i \leq d} \left\langle \sum_{t=0}^T u_t, e_i \right\rangle = \max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x_t \rangle.$$

Therefore, upper bounds on the quantity

$$\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x \rangle - \sum_{t=0}^T \langle u_t, x_t \rangle = \max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - x_t \rangle, \quad (3.2)$$

which we will also call the *regret*, will yield the same bound on the expected regret from (3.1).

A common assumption is that the sequence of payoff vectors $(u_t)_{t \geq 0}$ is bounded. In that case, we will see that there exists algorithms that

¹This interpretation holds only if we assume that the choice of the payoff vectors $(u_t)_{t \geq 0}$ by Nature does not depend on the choices of the Decision Maker. Nevertheless, all guarantees on the regret will still hold, even in the case where Nature does react to the choices of the Decision Maker; only the interpretation will not stand.

guarantee that above (cumulative) regret grows at most in \sqrt{T} , in other words, the *average* regret is bounded from above by a quantity that vanishes as $1/\sqrt{T}$. This has the following interpretation called *prediction with expert advice*: there are d experts, and at each step $t \geq 0$, the Decision Maker has to choose one of the experts and follow his advice, and obtains the corresponding payoff. Then, as will be proved below, there exists algorithms such that the Decision Maker is guaranteed to perform *as well as the best expert* (asymptotically and in average).

Another important question will be the optimal dependence of the regret bound in d , in the case e.g. where the payoff vectors are assumed to be in $[0, 1]^d$.

Example 3.1.1 (Follow the leader). The simplest algorithm we may think of is called *follow the leader*, and picks the decision $i \in \{1, \dots, d\}$ which would have yielded the highest cumulative payoff on the previous steps:

$$i_{t+1} = \arg \max_{1 \leq i \leq d} \sum_{s=1}^t u_{s,i}, \quad t \geq 0,$$

which can be equivalently written

$$x_{t+1} = \arg \max_{x \in \Delta_d} \left\langle \sum_{s=1}^t u_s, x \right\rangle, \quad t \geq 0. \quad (3.3)$$

Unfortunately, this algorithm is too simple and it is easy to find a bounded sequence for which the regret grows linearly, e.g. for $d = 2$,

$$u_0 = \begin{pmatrix} 1/2 \\ 0 \end{pmatrix}, \quad u_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad u_3 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \dots$$

yields for all $T \geq 1$,

$$\max_{1 \leq i \leq d} \sum_{t=0}^T u_{t,i} - \sum_{t=0}^T u_{t,i_t} \geq \frac{T-1}{2} - \frac{1}{2}.$$

Intuitively, the issue with the follow the leader algorithm is that it follows previous data too closely, so that there exists a payoff vector for the next step for which the decision of the algorithm is the worst. This is a kind of *overfitting*. To address the issue, one possible approach is to *regularize* the quantity that is maximized in (4.2), which will lead to the *dual averaging* and *follow the regularized leader* algorithms below.

We now make the connection with online learning and optimization by presenting two successive extensions of the above framework.

Online linear optimization The quantity (3.2) inspires the following natural extension. Let $\mathcal{X} \subset \mathbb{R}^d$ be a nonempty closed convex set, and $\mathcal{U} \subset \mathbb{R}^d$ any nonempty set. At each step $t \geq 0$,

- the Decision Maker chooses $x_t \in \mathcal{X}$,
- Nature chooses and reveals $u_t \in \mathcal{U}$,
- the Decision Maker obtains payoff $\langle u_t, x_t \rangle$.

In the case where \mathcal{X} is bounded, the natural definition of the regret is

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T \langle u_t, x - x_t \rangle.$$

When \mathcal{X} is unbounded, it will be possible to guarantee upper bounds on

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle,$$

that depend on the *comparison point* $x \in \mathcal{X}$. In the case where the set \mathcal{U} of payoff vectors is bounded, typical regret bounds also grow as \sqrt{T} , as will be established below in Sections 3.2 and 3.3.

Online convex optimization We now further generalize by considering convex loss functions, which is motivated by e.g. online learning. Let $\mathcal{X} \subset \mathbb{R}^d$ be a nonempty closed convex set and \mathcal{L} a nonempty set of convex function $\ell : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ such that $\mathcal{X} \subset \text{dom } \ell$. At each step $t \geq 0$,

- the Decision Maker chooses $x_t \in \mathcal{X}$,
- Nature chooses and reveals $\ell_t \in \mathcal{L}$,
- the Decision Maker incurs loss $\ell_t(x_t)$.

The corresponding definition for the regret is

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)).$$

In the case where the loss functions have some *curvature*, the latter can be leveraged to achieve regret bounds that grow strictly slower than \sqrt{T} .

The case where the loss function is constant boils down to convex optimization.

Example 3.1.2 (Online linear regression). By slightly modifying the above problem, the Decision Maker can observe some contextual information z_t before choosing $x_t \in \mathcal{X}$. Online versions of supervised learning problem can then be considered with Nature choosing loss function of the form e.g.

$$\ell(x) = (\langle z, x \rangle - y)^2, \quad z \in \mathbb{R}^d, y \in \mathbb{R}.$$

3.2 Dual averaging

We define and analyze the dual averaging family of algorithm which can be seen as a *regularized* version of the follow the leader algorithm from Example (4.2). It is a special case of UMD iterates where for all $t \geq 0$, the regularizers satisfy $h_{t+1/2} = h_{t+1}$ and where the next dual point y_{t+1} is uniquely defined as $y_{t+1} = y_t + u_t$.

We obtain in Proposition 3.2.5 below a regret bound in the context of online linear optimization, which will be applied and transposed to numerous problems.

Definition 3.2.1. Let $(h_t)_{t \geq 0}$ be a sequence of regularizers on \mathcal{X} , $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , and $y_0 \in \mathbb{R}^d$. The associated sequence $(x_t)_{t \geq 0}$ of *dual averaging* (DA) iterates is defined for $t \geq 0$ as

$$x_t = \nabla h_t^*(y_t) \quad \text{and} \quad y_{t+1} = y_t + u_t.$$

Remark 3.2.2. The above definition can be equivalently written

$$\begin{aligned} x_{t+1} &= \arg \max_{x \in \mathbb{R}^d} \left\{ \left\langle y_0 + \sum_{s=0}^t u_s, x \right\rangle - h_t(x) \right\} \\ &= \arg \max_{x \in \mathcal{X}} \{ \langle u_t, x \rangle - D_h(x, x_t; y_t) \}. \end{aligned}$$

In the case $y_0 = 0$, the above second expression gives the following interpretation: x_{t+1} is the maximizer not of the past cumulative payoff function $x \mapsto \langle \sum_{s=0}^t u_s, x \rangle$ as in the *follow the leader* algorithm (see Section 3.1), but of a *regularized* version. For this reason, this algorithm is sometimes called *follow the regularized leader*. We will use this name below to designate a more general algorithm in the context of regret minimization with convex losses.

Example 3.2.3 (Lazy Mirror Descent). Let $y_0 \in \mathbb{R}^d$ and denote $\Pi_{\mathcal{X}}$ the Euclidean projection onto \mathcal{X} . For a given sequence $(u_t)_{t \geq 0}$, $x_0 = \Pi_{\mathcal{X}}(y_0)$ and

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t + u_t), \quad t \geq 0$$

corresponds to dual averaging iterates with constant Euclidean regularizer $h = \frac{1}{2} \|\cdot\|_2^2 + I_{\mathcal{X}}$ (see Proposition 2.2.5 and Example 2.3.6).

Example 3.2.4 (Exponential weights algorithm). For a given sequence $(u_t)_{t \geq 0}$, consider

$$x_{t+1} = \left(\frac{\exp(\sum_{s=0}^t u_{s,i})}{\sum_{j=1}^d \exp(\sum_{s=0}^t u_{s,j})} \right)_{1 \leq j \leq d}.$$

These correspond to dual averaging iterates with constant entropic regularizer h_{ent} (see Proposition 2.2.6)

The following statement gives a regret bound for online linear optimization, which will also be applied and transposed to numerous problems.

Proposition 3.2.5 (Regret bounds for DA). *Let $(x_t)_{t \geq 0}$ and $(y_t)_{t \geq 0}$ be defined as in Definition 3.2.1, and $x \in \bigcap_{t \geq 0} \text{dom } h_t$. Then,*

(i) *$((x_t, y_t))_{t \geq 0}$ is a sequence of UMD iterates associated with regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ and dual increments $(u_t)_{t \geq 0}$, where $h_{t+1/2} := h_{t+1}$ for all $t \geq 0$;*

(ii) *for all $T \geq 0$,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq D_0 + (h_{T+1}(x) - h_0(x)) + \sum_{t=0}^T D_t^*,$$

where

$$\begin{aligned} D_0 &= h_0(x) - h_0(x_0) - \langle y_0, x - x_0 \rangle \\ D_t^* &= h_{t+1}^*(y_t + u_t) - h_t^*(y_t) - \langle u_t, x_t \rangle, \quad t \geq 0; \end{aligned}$$

(iii) *if for $t \geq 0$, $h_{t+1} \geq h_t$ and h_t is K_t -strongly convex for some norm $\|\cdot\|$, then D_t^* is bounded as*

$$D_t^* \leq \frac{1}{2K_t} \|u_t\|_*^2.$$

Proof. (i) holds because the conditions from Definition 2.3.1 are trivially satisfied. (ii) and (iii) easily follow from Lemma 2.5.2. \square

Dual averaging with nonincreasing parameter Let h be a regularizer on \mathcal{X} , $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , $y_0 = 0$ and $(\eta_t)_{t \geq 0}$ a positive nonincreasing sequence. Consider iterates $(x_t)_{t \geq 0}$ defined for $t \geq 0$ as

$$x_t = \nabla h^*(\eta_t y_t) \quad \text{and} \quad y_{t+1} = y_t + u_t. \quad (3.4)$$

Then, $((x_t, y_t))_{t \geq 0}$ are UMD iterates associated with dual increments $(u_t)_{t \geq 0}$ and regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ where

$$h_t(x) = \frac{h(x) - \min h}{\eta_t}, \quad t \geq 0, \quad (3.5)$$

and $h_{t+1/2} = h_{t+1}$ for $t \geq 1$. The definition of UMD iterates is invariant when constants are added to regularizers, so it would be equivalent to simply consider $h_t = h/\eta_t$ but the above regularizers have the advantage of ensuring $h_{t+1} \geq h_t$ and therefore makes the analysis simpler.

Proposition 3.2.6 (Regret bounds for DA with time-dependent parameters). *Consider the iterates defined in (3.4).*

(i) For all $T \geq 0$ and $x \in \text{dom } h$,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{h(x) - \min h}{\eta_T} + \sum_{t=0}^T \frac{D_{h^*}(\eta_t(y_t + u_t), \eta_t y_t)}{\eta_t}.$$

(ii) Moreover, if h is K -strongly convex for some norm $\|\cdot\|$, for all $T \geq 0$,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{h(x) - \min h}{\eta_T} + \frac{1}{2K} \sum_{t=0}^T \eta_t \|u_t\|_*^2.$$

(iii) Moreover, if there exists $L > 0$ such that $\|u_t\|_* \leq L$, then the choice $\eta_t = \eta\sqrt{2K}/(L\sqrt{t+1})$ for some $\eta > 0$ yields

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \left(\frac{h(x) - \min h}{\eta} + \eta \right) L \sqrt{\frac{T+1}{2K}}.$$

(iv) Moreover, if $\sup_{x \in \mathcal{X}} h(x) < +\infty$, then $\eta = \sqrt{\max_{\mathcal{X}} h - \min h}$ yields

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T \langle u_t, x - x_t \rangle \leq L \sqrt{\frac{2(\max_{\mathcal{X}} h - \min h)(T+1)}{K}}.$$

Proof. (i) Applying Proposition 3.2.5, using $y_0 = 0$, and simplifying gives

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{h(x) - \min h}{\eta_{T+1}} + \sum_{t=0}^T (h_{t+1}^*(y_t + u_t) - h_t^*(y_t) - \langle u_t, x_t \rangle).$$

For a given $T \geq 0$, iterates x_0, \dots, x_T do not depend on η_{T+1} , therefore, Proposition 3.2.5 can be applied for $\eta_{T+1} = \eta_T$. Besides, because $h_{t+1} \geq h_t$ by definition in (3.5), $h_{t+1}^*(y_t + u_t) \leq h_t^*(y_t + u_t)$, hence the result.

(ii) follows from Proposition 1.6.4, (iii) and (iv) are immediate consequences. \square

Remark 3.2.7. According to the above statements (iii) and (iv) give that the average regret is minimized at speed $1/\sqrt{T}$, for all $T \geq 0$, without prior knowledge of the latter: the algorithm and the guarantee are said to be *horizon-free*. In the case of a bounded domain such that $\sup_{\mathcal{X}} h < +\infty$, the bound (iv) does not depend on the comparison point x . When \mathcal{X} is unbounded, or when $\sup_{\mathcal{X}} h = +\infty$, the bound (iii) depend on the comparison point x , but still guarantees a horizon-free average regret bound of order $1/\sqrt{T}$, which is an important feature of dual averaging.

3.3 Online mirror descent

We define and analyze the the online mirror descent family of algorithms, which is an extension of the online projected gradient descent:

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t + u_t) = \arg \max_{x \in \mathcal{X}} \left\{ \langle u_t, x \rangle - \frac{1}{2} \|x - x_t\|_2^2 \right\},$$

where the above Euclidean distance is replaced by the Bregman divergence associated with a differentiable function H . This yields a special case of UMD iterates where for all $t \geq 0$, $h_t = h_{t+1/2}$ and where dual point y_t is uniquely defined as $y_t = \nabla H_t(x_t)$.

Proposition 3.3.13 below gives a regret bound in the context of online linear optimization, and will also be applied and transposed to various problems.

Definition 3.3.1. Let $H : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$. Denote $\mathcal{D}_H := \text{int dom } H$. H is a *mirror map* compatible with \mathcal{X} if

- (i) H is lower semi-continuous, strictly convex, and differentiable on \mathcal{D}_H ,
- (ii) the gradient of H takes all possible values, i.e. $\nabla H(\mathcal{D}_H) = \mathbb{R}^d$,
- (iii) $\mathcal{X} \subset \text{cl } \mathcal{D}_H$,
- (iv) $\mathcal{X} \cap \mathcal{D}_H \neq \emptyset$.

Example 3.3.2 (ℓ^p norms). For $p > 1$, $x \mapsto \frac{1}{2} \|x\|_p^2$ is a mirror map compatible with all nonempty closed convex sets.

Example 3.3.3 (Generalized negative entropy). The generalized negative entropy on the closed positive orthant, defined as

$$H(x) = \begin{cases} \sum_{i=1}^d x_i \log x_i & \text{if } x \in \mathbb{R}_+^d \\ +\infty & \text{otherwise,} \end{cases}$$

with convention $0 \cdot \log 0 = 0$, is a mirror map compatible with all nonempty closed convex subsets of \mathbb{R}_+^d (e.g. the simplex).

Example 3.3.4 (Log barrier). The log barrier on the open positive orthant, defined as

$$H(x) = \begin{cases} -\sum_{i=1}^d \log x_i & \text{if } x \in (\mathbb{R}_+^*)^d, \\ +\infty & \text{otherwise,} \end{cases}$$

is a mirror map compatible with all nonempty closed convex subsets of \mathbb{R}_+^d (e.g. the simplex), although its domain is the open positive orthant $(\mathbb{R}_+^*)^d$.

Proposition 3.3.5. Let H be a mirror map compatible with \mathcal{X} , H^* its Legendre–Fenchel transform. Then,

$$(i) \text{ dom } H^* = \mathbb{R}^d,$$

$$(ii) H^* \text{ is differentiable on } \mathbb{R}^d,$$

$$(iii) \nabla H^*(\mathbb{R}^d) = \mathcal{D}_H,$$

$$(iv) \text{ For } x \in \mathcal{D}_H \text{ and } y \in \mathbb{R}^d, \nabla F^*(\nabla F(x)) = x \text{ and } \nabla F(\nabla F^*(y)) = y.$$

Proof. Let $y_t \in \mathbb{R}^n$. By property (ii) from Definition 3.3.1, there exists $x_0 \in \mathcal{D}_H$ such that $\nabla H(x_0) = y$. Therefore, function $\varphi_y : x \mapsto \langle y|x \rangle - H(x)$ is differentiable at x_0 and $\nabla \varphi_y(x_0) = 0$. Moreover, φ_y is strictly concave as a consequence of property (i) from Definition 3.3.1. Therefore, x_0 is the unique maximizer of φ_y and:

$$H^*(y) = \max_{x \in \mathbb{R}^n} \{\langle y|x \rangle - H(x)\} < +\infty,$$

which proves property (i).

Besides, we have

$$x_0 \in \partial H^*(y) \iff y = \nabla H(x_0) \iff x_0 \text{ minimizer of } \phi_y, \quad (3.6)$$

where the first equivalence comes from Proposition 1.4.6. Point x_0 being the unique maximizer of φ_y , we have that $\partial H^*(y)$ is a singleton. In other words, H^* is differentiable in y and

$$\nabla H^*(y) = x_0 \in \mathcal{D}_H. \quad (3.7)$$

First, the above (3.7) proves property (ii). Second, this equality combined with the equality from (3.6) gives the second identity from property (iv). Third, this proves that $\nabla H^*(\mathbb{R}^n) \subset \mathcal{D}_H$.

It remains to prove the reverse inclusion to get property (iii). Let $x \in \mathcal{D}_H$. By property (i) from Definition 3.3.1, H is differentiable in x . Consider

$$y := \nabla H(x), \quad (3.8)$$

and all the above holds with this special point y . In particular, $x_0 = x$ by uniqueness of x_0 . Therefore (3.7) gives

$$\nabla H^*(y) = x, \quad (3.9)$$

and this proves $\nabla H^*(\mathbb{R}^n) \supset \mathcal{D}_H$ and thus property (iii). Combining (3.8) and (3.9) gives the first identity from property (iv). \square

Proposition 3.3.6 (OMD iteration). *Let H be a mirror map compatible with \mathcal{X} , $x \in \mathcal{D}_H$, $u \in \mathbb{R}^d$ and consider*

$$x' = \arg \max_{x'' \in \mathcal{X}} \{\langle u, x'' \rangle - D_H(x'', x)\}.$$

Then, $x' \in \mathcal{X} \cap \mathcal{D}_H$ and can also be written

$$\begin{aligned} x' &= \arg \max_{x'' \in \mathcal{X}} \{ \langle \nabla H(x) + u, x'' \rangle - H(x'') \} \\ &= \arg \min_{x'' \in \mathcal{X}} D_H(x'', \nabla H^*(\nabla H(x) + u)). \end{aligned}$$

Proof. The proof that x' is well-defined and belongs to \mathcal{D}_H is given in [JKM23]. Besides,

$$\begin{aligned} x' &= \arg \max_{x'' \in \mathcal{X}} \{ \langle u, x'' \rangle - D_H(x'', x) \} \\ &= \arg \max_{x'' \in \mathcal{X}} \{ \langle u, x'' \rangle - H(x'') + \langle \nabla H(x), x'' - x \rangle \} \\ &= \arg \max_{x'' \in \mathcal{X}} \{ \langle \nabla H(x) + u, x'' \rangle - H(x'') \} \\ &= \arg \max_{x'' \in \mathcal{X}} \{ -H(x'') + H(\nabla H^*(\nabla H(x) + u)) + \langle \nabla H(x) + u, x'' - x \rangle \} \\ &= \arg \min_{x'' \in \mathcal{X}} D_H(x'', \nabla H^*(\nabla H(x) + u)). \end{aligned}$$

□

Definition 3.3.7. A sequence $(H_t)_{t \geq 0}$ of mirror maps has *nondecreasing domains* if $\text{dom } H_t \subset \text{dom } H_{t+1}$ for all $t \geq 0$.

Definition 3.3.8. Let $(H_t)_{t \geq 0}$ be a sequence of mirror maps compatible with \mathcal{X} with nondecreasing domains and $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d . A sequence $(x_t)_{t \geq 0}$ in \mathbb{R}^d is an sequence of *online mirror descent (OMD)* iterates on \mathcal{X} associated with mirror maps $(H_t)_{t \geq 0}$ and *dual increments* $(u_t)_{t \geq 0}$ if $x_0 \in \mathcal{X} \cap \text{int dom } H_0$ and

$$x_{t+1} = \arg \max_{x \in \mathcal{X}} \{ \langle u_t, x \rangle - D_{H_t}(x, x_t) \}, \quad t \geq 0.$$

Remark 3.3.9. The above iterates are well-defined. Indeed, we first note that $x \in \mathcal{D}_{H_0}$, and then by induction, as soon as $x_t \in \mathcal{D}_{H_t}$, Proposition 3.3.6 ensures that x_{t+1} is well-defined and unique, and that it belongs to \mathcal{D}_{H_t} . Because the mirror maps have nondecreasing domains, x_{t+1} also belongs to $\mathcal{D}_{H_{t+1}}$.

Example 3.3.10 (Online (projected) gradient descent). Denote $\Pi_{\mathcal{X}}$ the Euclidean projection onto \mathcal{X} . Let $y_0 \in \mathbb{R}^d$ and a sequence $(u_t)_{t \geq 0}$ in \mathbb{R}^d . Then, $x_0 = \Pi_{\mathcal{X}}(y_0)$ and

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t + u_t), \quad t \geq 0,$$

are OMD iterates on \mathcal{X} associated with constant Euclidean mirror map $H = \frac{1}{2} \|\cdot\|_2^2$.

Example 3.3.11 (Exponential weights algorithm). The exponential weights algorithm from Example 3.2.4 gives OMD iterates on the simplex Δ_d associated with generalized entropic mirror map from Example 3.3.3.

The following proposition indicates the regularizers to consider for making the connection with the definition of UMD iterates.

Proposition 3.3.12. *Let H be a mirror map compatible with \mathcal{X} . Then $h = H + I_{\mathcal{X}}$ is a regularizer on \mathcal{X} and for all $x \in \mathcal{D}_H$, $\nabla H(x) \in \partial h(x)$.*

Proof. See [JKM23]. \square

The following statement gives a regret bound for online linear optimization, and will also be applied and transposed to numerous problems.

Proposition 3.3.13 (Regret bounds for OMD). *Let $(H_t)_{t \geq 0}$ be a sequence of mirror maps compatible with \mathcal{X} with nondecreasing domains, $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , $(x_t)_{t \geq 0}$ associated OMD iterates in \mathcal{X} and $x \in \mathcal{X} \cap \text{dom } H_0$. Then,*

(i) *$((x_t, \nabla H_t(x_t)))_{t \geq 0}$ is a sequence of UMD iterates associated with regularizers $(h_t)_{t \geq 0}$ defined as*

$$h_t = h_{t+1/2} = H_t + I_{\mathcal{X}}, \quad t \in \mathbb{N},$$

and dual increments $(u_t)_{t \geq 0}$;

(ii) *for all $T \geq 0$,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq D_{H_0}(x, x_0) - D_{H_{T+1}}(x, x_{T+1}) + \sum_{t=0}^T (\tilde{D}_{t+1/2}^{\Delta} + \tilde{D}_t^*),$$

where

$$\begin{aligned} \tilde{D}_{t+1/2}^{\Delta} &= D_{H_{t+1}-H_t}(x, x_{t+1}), \\ \tilde{D}_t^* &= D_{H_t^*}(\nabla H_t(x_t) + u_t, \nabla H_t(x_t)); \end{aligned}$$

(iii) *if for $t \geq 0$, H_t is K_t -strongly convex for some norm $\|\cdot\|$, then \tilde{D}_t^* is bounded as*

$$\tilde{D}_t^* \leq \frac{1}{2K_t} \|u_t\|_*^2.$$

Proof. (i) $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ are indeed regularizers thanks to Proposition 3.3.12. Let us prove that $((x_t, \nabla H_t(x_t)))_{t \geq 0}$ satisfy the definition of UMD iterates. Let $t \geq 0$ and $x' \in \mathbb{R}^d$. By definition of h_t , $h_t \geq H_t$. Moreover, $h_t(x_t) = H(x_t)$ because $x_t \in \mathcal{D}_{H_t}$ by Remark 3.3.9. Then,

$$h_t(x') - h_t(x_t) \geq H_t(x') - H_t(x_t) \geq \langle \nabla H_t(x_t), x - x_t \rangle,$$

which means $\nabla F_t(x_t) \in \partial h_t(x_t)$, and the first condition from Definition 2.3.1 is satisfied. Besides, using Proposition 3.3.6 and the definition of $h_{t+1/2}$,

$$\begin{aligned} x_{t+1} &= \arg \max_{x \in \mathcal{X}} \{ \langle \nabla H_t(x_t) + u_t, x \rangle - H_t(x) \} \\ &= \arg \max_{x \in \mathbb{R}^d} \{ \langle \nabla H_t(x_t) + u_t, x \rangle - h_{t+1/2}(x) \} = \nabla h_{t+1/2}^*(\nabla H_t(x_t) + u_t), \end{aligned}$$

which proves the second condition, and $((x_t, \nabla H_t(x_t)))_{t \geq 0}$ is indeed a sequence of UMD iterates associated with $(H_t)_{t \geq 0}$ and $(u_t)_{t \geq 0}$.

(ii) For all $t \in \frac{1}{2}\mathbb{N}$, the definition of h_t imply that $\text{dom } h_t = \mathcal{X} \cap \text{dom } H_t$. Because $(H_t)_{t \geq 0}$ has nondecreasing domains,

$$\begin{aligned} x \in \mathcal{X} \cap \text{dom } H_0 &= \mathcal{X} \cap \bigcap_{t \in \mathbb{N}} \text{dom } H_t = \bigcap_{t \in \mathbb{N}} (\mathcal{X} \cap \text{dom } H_t) \\ &= \bigcap_{t \in \mathbb{N}} \text{dom } h_t = \bigcap_{t \in \frac{1}{2}\mathbb{N}} \text{dom } h_t. \end{aligned}$$

Therefore, Lemma 2.5.2 can be applied with x and with notation therein, we get for all $T \geq 0$,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle = D_0 - D_{T+1} + \sum_{t=0}^T (D_{t+1/2}^\Delta + D_t^*).$$

First, because $x, x_0 \in \mathcal{X} \cap \text{dom } H_0$,

$$\begin{aligned} D_0 &= h_0(x) - h_0(x_0) - \langle \nabla H_0(x_0), x - x_0 \rangle \\ &= H_0(x) - H_0(x_0) - \langle \nabla H_0(x_0), x - x_0 \rangle \\ &= D_{H_0}(x, x_0). \end{aligned}$$

Similarly, x belongs to $\mathcal{X} \cap \text{dom } H_{T+1}$ and so does x_{T+1} as a consequence of $(x_t)_{t \geq 0}$ being OMD iterates (see Remark 3.3.9) and $D_{T+1} = D_{H_{T+1}}(x, x_{T+1})$.

Let us now bound the two remaining Bregman divergences. For $t \geq 0$, denote $y_t = \nabla H_t(x_t)$, then Proposition 3.3.6 ensures that

$$x_{t+1} = \arg \max_{x \in \mathcal{X}} \{ \langle y_t + u_t, x \rangle - H_t(x) \},$$

in other words x_{t+1} is a minimizer of a differentiable convex function over \mathcal{X} . Therefore, Proposition 1.2.10 gives the following variational characterization:

$$\langle \nabla H_t(x_{t+1}) - y_t - u_t, x' - x_{t+1} \rangle \geq 0, \quad x' \in \mathcal{X}.$$

We use it for $x' = x$ to bound $D_{t+1/2}^\Delta$ as follows.

$$\begin{aligned}
D_{t+1/2}^\Delta &= \Delta h_{t+1/2}(x) - \Delta h_{t+1/2}(x_{t+1}) \\
&\quad - \langle y_{t+1} - y_t - u_t, x - x_{t+1} \rangle \\
&\leq H_{t+1}(x) - H_t(x) - H_{t+1}(x_{t+1}) + H_t(x_{t+1}) \\
&\quad - \langle \nabla H_{t+1}(x_{t+1}) - \nabla H_t(x_{t+1}), x - x_{t+1} \rangle \\
&= D_{H_{t+1}-H_t}(x, x_{t+1}) \\
&= \tilde{D}_{t+1/2}^\Delta.
\end{aligned}$$

We now turn to D_t^* . Because $h_t \geq H_t$ by definition of h_t , the reverse inequality holds for the Legendre–Fenchel transform, and in particular, $h_t^*(y_t + u_t) \leq H_t^*(y_t + u_t)$. Besides, it holds that

$$x_t = \nabla H_t^*(y_t) \quad \text{and} \quad x_t = \nabla h_t^*(y_t),$$

where the first equality comes from Proposition 3.3.5, whereas the second is obtained combining the fact that $\nabla H_t(x_t) = y_t \in \partial h_t(x_t)$ (proved above) with Proposition 1.4.6. The latter proposition also gives a characterization of both above gradients as maximizers of concave functions, which yields

$$\begin{aligned}
h_t^*(y_t) &= \max_{x \in \mathbb{R}^d} \{ \langle y_t, x \rangle - h_t(x) \} = \langle y_t, x_t \rangle - h_t(x_t) \\
&= \langle y_t, x_t \rangle - H_t(x_t) = \max_{x \in \mathbb{R}^d} \{ \langle y_t, x \rangle - H_t(x) \} \\
&= H_t^*(y_t).
\end{aligned}$$

Hence

$$\begin{aligned}
D_t^* &= h_{t+1/2}^*(y_t + u_t) - h_t^*(y_t) - \langle u_t, x_t \rangle \leq H_t^*(y_t + u_t) - H_t^*(y_t) - \langle u_t, x_t \rangle \\
&= D_{H_t^*}(y_t + u_t, y_t).
\end{aligned}$$

□

OMD with nonincreasing step-sizes An important special case of OMD iterates is the following. Let H be a mirror map compatible with \mathcal{X} , $(\gamma_t)_{t \geq 0}$ a positive and nonincreasing sequence, and $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d . Then, associated OMD iterates on \mathcal{X} with time-dependent *step-sizes* (or *learning rate*) are defined as

$$x_{t+1} = \arg \max_{x \in \mathcal{X}} \{ \langle \gamma_t u_t, x \rangle - D_H(x, x_t) \} \quad (3.10)$$

$$= \arg \max_{x \in \mathcal{X}} \{ \langle \nabla H(x_t) + \gamma_t u_t, x \rangle - H(x) \}. \quad (3.11)$$

This corresponds to OMD iterates from Definition 3.3.8 with time-dependent mirror maps $(H_t)_{t \geq 0}$ defined as $H_t = \gamma_t^{-1} H$ for all $t \geq 0$, which is indeed a sequence of mirror maps compatible with \mathcal{X} with nondecreasing domains.

Proposition 3.3.14 (Regret bounds for OMD with time-dependent step-sizes). *Consider the OMD iterates with nonincreasing step-sizes defined in (3.10).*

(i) *For all $T \geq 0$ and $x \in \mathcal{X} \cap \text{dom } H$,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{\max_{0 \leq t \leq T} D_H(x, x_t)}{\gamma_T} + \sum_{t=0}^T \tilde{D}_t^*,$$

where $\tilde{D}_t^* = \gamma_t^{-1} D_{H^*}(\nabla H(x_t) + \gamma_t u_t, \nabla H_t(x_t))$.

(ii) *Moreover, if H is K -strongly convex with respect to some norm $\|\cdot\|$, for all $T \geq 0$,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{\max_{0 \leq t \leq T} D_H(x, x_t)}{\gamma_T} + \frac{1}{2K} \sum_{t=0}^T \gamma_t \|u_t\|_*^2.$$

(iii) *Moreover, if there exists known constants $R > 0$ and $L > 0$, such that for all $t \geq 0$, $\max_{x \in \mathcal{X}} D_H(x, x_t) \leq R^2$ and $\|u_t\|_* \leq L$, then step-sizes $\gamma_t = R\sqrt{K}/(L\sqrt{t+1})$ yield for all $T \geq 0$,*

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T \langle u_t, x - x_t \rangle \leq RL \sqrt{\frac{2(T+1)}{K}}.$$

Proof. (i) Applying Proposition 3.3.13,

$$\begin{aligned} \sum_{t=0}^T \langle u_t, x - x_t \rangle &= \frac{D_H(x, x_0)}{\gamma_0} - \frac{D_H(x, x_{T+1})}{\gamma_{T+1}} \\ &\quad + \sum_{t=0}^T \left(\frac{1}{\gamma_{t+1}} - \frac{1}{\gamma_t} \right) D_H(x, x_{t+1}) + \sum_{t=0}^T \tilde{D}_t^* \\ &= \frac{D_H(x, x_0)}{\gamma_0} + \sum_{t=0}^{T-1} \left(\frac{1}{\gamma_{t+1}} - \frac{1}{\gamma_t} \right) D_H(x, x_{t+1}) \\ &\quad - \frac{D_H(x, x_{T+1})}{\gamma_T} + \sum_{t=0}^T \tilde{D}_t^* \\ &\leq \left(\max_{1 \leq t \leq T} D_H(x, x_t) \right) \left(\frac{1}{\gamma_0} + \sum_{t=0}^{T-1} \left(\frac{1}{\gamma_{t+1}} - \frac{1}{\gamma_t} \right) \right) + \sum_{t=0}^T \tilde{D}_t^* \\ &= \frac{\max_{1 \leq t \leq T} D_H(x, x_t)}{\gamma_T} + \sum_{t=0}^T \tilde{D}_t^*. \end{aligned}$$

Besides, for all $t \geq 0$ and $y \in \mathbb{R}^d$,

$$H_t^*(y) = \max_{x \in \mathbb{R}^d} \left\{ \langle y, x \rangle - \frac{H(x)}{\gamma_t} \right\} = \frac{1}{\gamma_t} \max_{x \in \mathbb{R}^d} \{ \langle \gamma_t y, x \rangle - H(x) \},$$

which yields

$$\tilde{D}_t^* = D_{H_t^*}(\nabla H_t(x_t) + u_t, \nabla H_t(x_t)) = \frac{D_{H^*}(\nabla H(x_t) + \gamma_t u_t, \nabla H(x_t))}{\gamma_t}.$$

(ii) and (iii) follow immediately. \square

Remark 3.3.15. Even for a fixed point $x \in \mathcal{X} \cap \text{dom } H$, unless there is a known bound on $\max_{t \geq 0} D_H(x, x_t)$, for instance in the case where \mathcal{X} is bounded, there is no known sublinear regret bound for OMD. This is contrast with DA which, with the right nonincreasing parameters from Proposition 3.2.5, guarantees a regret bound that grows as \sqrt{T} with time, as soon as a bound on the norms of vectors $(u_t)_{t \geq 0}$ is known.

Corollary 3.3.16. *Consider the special case of online gradient descent iterations*

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t + \gamma_t u_t), \quad t \geq 0.$$

Then, for all $T \geq 0$ and $x \in \mathcal{X}$,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{\max_{0 \leq t \leq T} \|x - x_t\|_2^2}{2\gamma_T} + \sum_{t=0}^T \frac{\gamma_t \|u_t\|_2^2}{2}.$$

Proof. Combine Corollaries 3.3.16 and 1.6.8. \square

3.4 Finite action set

We recall the regret minimization problem where the Decision Maker chooses at each step an element in $\{1, \dots, d\}$ possibly at random, described in Section 3.1. At step $t \geq 0$,

- the Decision Maker chooses $x_t \in \Delta_d$,
- Nature chooses and reveals *payoff vector* $u_t \in [0, 1]^d$,
- i_t is a random element in $\{1, \dots, d\}$ drawn according to distribution x_t and revealed to the Decision Maker,
- the Decision Maker obtains payoff u_{t, i_t} .

As already discussed, the regret minimization in this setting can be reduced to an online linear optimization on the simplex Δ_d , as formalized in the following lemma.

Lemma 3.4.1. *For all $T \geq 0$,*

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right] = \mathbb{E} \left[\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - x_t \rangle \right].$$

Therefore, any upper bound on $\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - x_t \rangle$ will also be an upper bound on $\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right]$. A common assumption in this setting is that there is a known bound on the ℓ_∞ norm of the payoff vectors. We establish the important $\sqrt{T \log d}$ regret bound in this case.

Definition 3.4.2 (Exponential weights algorithm). Let $(u_t)_{t \geq 0}$ be a sequence in \mathbb{R}^d and $(\eta_t)_{t \geq 0}$ a positive sequence. The associated iterates of *exponential weights algorithm* (EW) iterates are defined in the simplex Δ_d as

$$x_t = \left(\frac{\exp \left(\eta_t \sum_{s=0}^{t-1} u_{s,i} \right)}{\sum_{j=1}^d \exp \left(\eta_t \sum_{s=0}^{t-1} u_{s,j} \right)} \right)_{1 \leq i \leq d}, \quad t \geq 0.$$

We recall the definition of the entropic regularizer on the simplex:

$$h_{\text{ent}}(x) = \begin{cases} \sum_{i=1}^d x_i \log x_i & \text{if } x \in \Delta_d, \text{ with convention } 0 \cdot \log 0 = 0 \\ +\infty & \text{otherwise.} \end{cases}$$

Proposition 3.4.3. (i) $h_{\text{ent}}^*(y) = \log \left(\sum_{i=1}^d \exp(y_i) \right)$ for all $y \in \mathbb{R}^d$.

$$(ii) \quad \nabla h_{\text{ent}}^*(y) = \left(\frac{\exp(y_i)}{\sum_{j=1}^d \exp(y_j)} \right)_{1 \leq i \leq d} \text{ for all } y \in \mathbb{R}^d,$$

$$(iii) \quad \max_{\Delta_d} h_{\text{ent}} - \min h_{\text{ent}} = \log d,$$

$$(iv) \quad h_{\text{ent}} \text{ is } 1\text{-strongly convex for } \|\cdot\|_1.$$

Proof. (iii) h_{ent} being convex, its maximum on Δ_d is attained at one of the extreme points. At each extreme point, the value of h_{ent} is zero. Therefore, $\max_{\Delta_d} h_{\text{ent}} = 0$. As for the minimum, h_{ent} being convex and symmetric with respect to the components x_i , its minimum is attained at the centroid $(1/d, \dots, 1/d)$ of the simplex Δ_d , where its value is $-\log d$. Therefore, $\min_{\Delta_d} h_{\text{ent}} = -\log d$, hence the result.

(iv) Consider $F : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by

$$F(x) = \begin{cases} \sum_{i=1}^d x_i (\log x_i - 1) & \text{if } x \in \mathbb{R}_+^d \\ +\infty & \text{otherwise.} \end{cases}$$

Let us prove that F is 1-strongly convex with respect to $\|\cdot\|_1$. By definition, the domain of F is \mathbb{R}_+^d . It is differentiable on the interior of the domain $(\mathbb{R}_+^*)^d$ and $\nabla F(x) = (\log x_i)_{1 \leq i \leq d}$ for $x \in (\mathbb{R}_+^*)^d$. Therefore, the norm of $\nabla F(x)$ goes to $+\infty$ when x converges to a boundary point of \mathbb{R}_+^d . [Roc70, Theorem 26.1] then assures that the subdifferential $\partial F(x)$ is empty as soon as $x \notin (\mathbb{R}_+^*)^d$. Therefore, condition (iii) from Proposition 1.6.4, which we aim at proving, can be written

$$\langle \nabla F(x') - \nabla F(x), x' - x \rangle \geq \|x' - x\|_1^2, \quad x, x' \in (\mathbb{R}_+^*)^d. \quad (3.12)$$

Let $x, x' \in (\mathbb{R}_+^*)^d$.

$$\langle \nabla F(x') - \nabla F(x), x' - x \rangle = \sum_{i=1}^d \log \frac{x'_i}{x_i} (x'_i - x_i).$$

A simple study of function shows that $(z-1) \log z - 2(z-1)^2/(z+1) \geq 0$ for $z \geq 0$. Applied with $z = x'_i/x_i$, this gives

$$\sum_{i=1}^d \log \frac{x'_i}{x_i} (x'_i - x_i) \geq \|x' - x\|_1^2,$$

and (4.3) is proved. F is therefore 1-strongly convex with respect to $\|\cdot\|_1$ and so is h_{ent} . \square

Proposition 3.4.4 (Regret bound for EW). *Let $(x_t)_{t \geq 0}$ be defined as in Definition 3.4.2. Then,*

(i) $(x_t)_{t \geq 0}$ is a sequence of DA iterates associated with regularizer h_{ent} , parameters $(\eta_t)_{t \geq 0}$ and dual increments $(u_t)_{t \geq 0}$.

(ii) Moreover, if $(\eta_t)_{t \geq 0}$ is nonincreasing, for all $T \geq 0$,

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right] \leq \frac{\log d}{\eta_T} + \sum_{t=0}^T \frac{D_{h_{\text{ent}}}^*(\eta_t(y_t + u_t), \eta_t y_t)}{\eta_t}.$$

(iii) If there exists $L > 0$ such that $\|u_t\|_\infty \leq L$ for all $t \geq 0$, and if $\eta_t = \sqrt{2 \log d} / (L\sqrt{t+1})$, then for all $T \geq 0$,

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right] \leq L\sqrt{2(\log d)(T+1)}.$$

Proof. (i) follows from Definition 3.4.2 and the expression of ∇h_{ent}^* from Proposition 3.4.3. (ii) and (iii) are then a simple application of Proposition 3.2.6 with the properties from Proposition 3.4.3 together with Lemma 3.4.1. \square

The above $\sqrt{T \log d}$ bound is known to be essentially unimprovable without further assumptions.

3.5 Multi-armed bandit

The multi-armed bandit problem is a variant of the regret minimization with a finite set of decisions where only the actual payoff is revealed to the Decision Maker. At step $t \geq 0$,

- the Decision Maker chooses $x_t \in \Delta_d$,
- Nature chooses $u_t \in \mathbb{R}^d$,
- i_t is drawn in $\{1, \dots, d\}$ according to x_t ,
- u_{t,i_t} is revealed to the Decision Maker.

We aim at obtaining guarantees on

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right].$$

Since the whole vector u_t is unknown to the Decision Maker, one possible approach is to construct an unbiased estimator of u_t as follows

$$\hat{u}_t = \left(\mathbb{1}_{\{i=i_t\}} \frac{u_{t,i_t}}{x_{t,i_t}} \right)_{1 \leq i \leq d}, \quad t \geq 0, \quad (3.13)$$

which indeed satisfies $\mathbb{E}[\hat{u}_t | x_t] = u_t$, and use it as a replacement to u_t in some online linear optimization algorithm on the simplex, e.g. the exponential weights algorithm. The resulting algorithm is called EXP3 (for exponential weights for exploration and exploitation) and is proved below to guarantee a regret bound of order $\sqrt{Td \log d}$, in the case where the payoff vectors are bounded with respect to $\|\cdot\|_\infty$.

Lemma 3.5.1. *For all $T \geq 0$,*

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right] \leq \mathbb{E} \left[\max_{x \in \Delta_d} \langle \hat{u}_t, x - x_t \rangle \right].$$

Proof. Using the fact that $\mathbb{E} \max \geq \max \mathbb{E}$,

$$\begin{aligned}
\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=1}^T \hat{u}_{t,i} - \sum_{t=1}^T \langle \hat{u}_t, x_t \rangle \right] &\geq \max_{1 \leq i \leq d} \mathbb{E} \left[\sum_{t=1}^T \hat{u}_{t,i} \right] - \mathbb{E} \left[\sum_{t=1}^T \langle \hat{u}_t, x_t \rangle \right] \\
&= \max_{1 \leq i \leq d} \mathbb{E} \left[\sum_{t=1}^T \mathbb{E} [\hat{u}_{t,i} | x_t] \right] \\
&\quad - \mathbb{E} \left[\sum_{t=1}^T \mathbb{E} [\langle \hat{u}_t, x_t \rangle | x_t] \right] \\
&= \max_{1 \leq i \leq d} \mathbb{E} \left[\sum_{t=1}^T u_{t,i} \right] - \mathbb{E} \left[\sum_{t=1}^T \langle u_t, x_t \rangle \right] \\
&= \max_{1 \leq i \leq d} \mathbb{E} \left[\sum_{t=1}^T u_{t,i} \right] - \mathbb{E} \left[\sum_{t=1}^T \mathbb{E} [u_{t,i_t} | x_t] \right] \\
&= \mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=1}^T u_{t,i} - \sum_{t=1}^T u_{t,i_t} \right],
\end{aligned}$$

here for the last equality, we used the fact that $u_{t,i}$ is deterministic to swap the maximum and the expectation. \square

For the analysis for EXP3, we need the following bound which provides a finer control over the Bregman divergence associated with h_{ent}^* .

Lemma 3.5.2. For $y \in \mathbb{R}^d$, $u \in \mathbb{R}_-^d$, $\eta > 0$ and $x = \nabla h_{\text{ent}}^*(\eta y)$,

$$D_{h_{\text{ent}}^*}(\eta(y+u), \eta y) \leq \frac{\eta^2}{2} \sum_{i=1}^d u_i^2 x_i.$$

Proof. Using the explicit expressions from Proposition 3.4.3

$$\begin{aligned}
D_{h_{\text{ent}}^*}(\eta(y+u), \eta y) &= h_{\text{ent}}^*(\eta(y+u)) - h_{\text{ent}}^*(\eta y) - \langle \nabla h_{\text{ent}}^*(\eta y) | \eta u \rangle \\
&= \log \left(\sum_{i=1}^d e^{\eta(y_i+u_i)} \right) - \log \left(\sum_{i=1}^d e^{\eta y_i} \right) - \eta \langle u, x \rangle \\
&= \log \left(\sum_{i=1}^d \frac{e^{\eta u_i} e^{\eta y_i}}{\sum_{j=1}^d e^{\eta y_j}} \right) - \eta \langle u, x \rangle \\
&= \log \left(\sum_{i=1}^d x_i e^{\eta u_i} \right) - \eta \langle u, x \rangle.
\end{aligned}$$

For $z \leq 0$, a simple differentiation proves that $e^z \leq 1 + z + \frac{z^2}{2}$. Therefore,

$$\begin{aligned} D_{h_{\text{ent}}^*}(\eta(y+u), \eta y) &\leq \log \left(\sum_{i=1}^d x_i \left(1 + \eta u_i + \frac{\eta^2 u_i^2}{2} \right) \right) - \eta \langle u, x \rangle \\ &= \log \left(1 + \eta \langle u, x \rangle + \frac{\eta^2}{2} \sum_{i=1}^d x_i u_i^2 \right) - \eta \langle u, x \rangle \\ &\leq \eta \langle u, x \rangle + \frac{\eta^2}{2} \sum_{i=1}^d x_i u_i^2 - \eta \langle u, x \rangle \\ &= \frac{\eta^2}{2} \sum_{i=1}^d x_i u_i^2, \end{aligned}$$

which gives the result. \square

Proposition 3.5.3 (Regret bound for EXP3). *Let $L > 0$ and $(u_t)_{t \geq 0}$ be a sequence in $[-L, 0]^d$. Let*

$$x_t = \left(\frac{\eta_t \sum_{s=0}^{t-1} \hat{u}_{s,i}}{\sum_{j=1}^d \eta_t \sum_{s=0}^{t-1} \hat{u}_{s,j}} \right)_{1 \leq i \leq d}, \quad t \geq 0,$$

where \hat{u}_t is defined as in (3.13) and $\eta_t = \sqrt{2(\log d)/(dL^2(t+1))}$. Then, for all $T \geq 0$,

$$\mathbb{E} \left[\max_{1 \leq i \leq d} \sum_{t=0}^T (u_{t,i} - u_{t,i_t}) \right] \leq L \sqrt{2(T+1)d \log d}.$$

Proof. Combine Proposition 3.4.4 with Lemmas 3.5.1 and 3.5.2. \square

Remark 3.5.4. An important limitation of the above result is that payoff vectors must belong to $[-L, 0]^d$. Although the regret is invariant when an additive constant is added to all components of a payoff vector, the construction of the estimator (3.13) however is not invariant by such a transformation. If payoff vectors are given in e.g. $[0, L]^d$, the same regret bound can be achieved by considering

$$\hat{u}_t = \left(\mathbb{1}_{\{i=i_t\}} \frac{u_{t,i_t} - L}{x_{t,i_t}} \right)_{1 \leq i \leq d}, \quad t \geq 0,$$

which are unbiased estimators of $u_t - L\mathbb{1}$.

Chapter 4

Online convex optimization

4.1 Introduction

Definition 4.1.1. $\ell : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is a *convex loss function* on \mathcal{X} if it is convex, lower semicontinuous, and has nonempty subdifferential on \mathcal{X} .

Remark 4.1.2. In particular, $\mathcal{X} \subset \text{dom } \ell$ according to Proposition 1.3.6.

We consider the online convex optimization problem. At each step $t \geq 0$,

- the Decision Maker chooses $x_t \in \mathcal{X}$,
- Nature chooses and reveals a convex loss function ℓ_t ,
- the Decision Maker incurs loss $\ell_t(x_t)$.

We aim at constructing algorithms which provide guarantees on the corresponding *regret*, defined as

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)), \quad T \geq 0, \quad x \in \mathcal{X}.$$

Example 4.1.3. *Online linear optimization* is a special case of online convex optimization where for all $t \geq 0$, loss functions are of the form $\ell_t(x) = -\langle u_t, x \rangle$ for some $u_t \in \mathbb{R}^d$. Then, the above definition of the regret coincide with the definition from Section 3.

Example 4.1.4. In the *online prediction with square loss* problem, for all $t \geq 0$, loss functions are of the form $\ell_t(x) = \frac{1}{2} \|x - z_t\|_2^2$ for some $z_t \in \mathbb{R}^d$.

Example 4.1.5. The *online portfolio optimization problem* models the sequential rebalancing of a portfolio between d assets. At each step $t \geq 0$, the Decision Maker chooses a distribution $x_t \in \Delta_d$ over the assets and changes the composition of its portfolio accordingly, so that the proportion (in value) corresponding to asset $i \in \{1, \dots, d\}$ is $x_{t,i}$. At the end of the step, the value

of each asset $i \in \{1, \dots, d\}$ is multiplied by a factor $r_{t,i} \in \mathbb{R}_+^*$ chosen by Nature, which corresponds to its performance. The value of the portfolio is then multiplied by $\langle r_t, x_t \rangle$. To fit into the online convex optimization framework, we consider the negative logarithm of the above quantity:

$$\ell_t(x) = -\log \langle r_t, x \rangle, \quad x \in \Delta_d,$$

so that the cumulative loss of the Decision Maker corresponds to the logarithm of the total variation ratio of the value of its portfolio.

4.2 Loss linearization

Online convex optimization can be reduced to an online linear optimization problem. Indeed, for $t \geq 0$, if $g_t \in \partial \ell_t(x_t)$, then by definition of a subgradient,

$$\ell_t(x_t) - \ell_t(x) \leq \langle g_t, x_t - x \rangle, \quad x \in \mathcal{X}. \quad (4.1)$$

Therefore, an upper bound on the regret corresponding to the online linear optimization problem with payoff vectors $(-g_t)_{t \geq 0}$ is also an upper bound on the initial regret corresponding to loss functions $(\ell_t)_{t \geq 0}$. The approach of using an online linear optimization algorithm with $(-g_t)_{t \geq 0}$ as payoff vectors is called *loss linearization*. The following guarantees are direct adaptations from Sections 3.2 and 3.3.

Corollary 4.2.1 (Online mirror descent for convex losses). *Let $K > 0$, $\|\cdot\|$ a norm in \mathbb{R}^d , H be a mirror map compatible with \mathcal{X} and K -strongly convex for $\|\cdot\|$, $(\gamma_t)_{t \geq 0}$ a positive and nonincreasing sequence, and $x_0 \in \mathcal{X} \cap \mathcal{D}_H$. Consider associated OMD iterates:*

$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \{ \langle \gamma_t g_t, x \rangle + D_H(x, x_t) \}, \quad t \geq 0.$$

Then for all $T \geq 0$ and $x \in \mathcal{X} \cap \text{dom } H$,

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{\max_{0 \leq t \leq T} D_H(x, x_t)}{\gamma_T} + \frac{1}{2K} \sum_{t=0}^T \gamma_t \|g_t\|_*^2.$$

Moreover, if there exists $R, L > 0$ such that for all $t \geq 0$, $\max_{x \in \mathcal{X}} D_H(x, x_t) \leq R^2$ and ℓ_t is L -Lipschitz for $\|\cdot\|$, then the choice

$$\gamma_t = \frac{R\sqrt{K}}{L\sqrt{t+1}}, \quad t \geq 0$$

guarantee for all $T \geq 0$,

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq RL \sqrt{\frac{2(T+1)}{K}}.$$

Corollary 4.2.2 (Dual averaging for convex losses). *Let $K > 0$, $\|\cdot\|$ a norm in \mathbb{R}^d , h a regularizer on \mathcal{X} which we assume K -strongly convex for $\|\cdot\|$, and $(\eta_t)_{t \geq 0}$ a positive and nonincreasing sequence. Consider associated DA iterates:*

$$x_t = \arg \min_{x \in \mathcal{X}} \left\{ \left\langle \eta_t \sum_{s=0}^{t-1} g_s, x \right\rangle + h(x) \right\}, \quad t \geq 0.$$

(i) *Then for all $T \geq 0$ and $x \in \text{dom } h$,*

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{h(x) - \min h}{\eta_T} + \frac{1}{2K} \sum_{t=0}^T \eta_t \|g_t\|_*^2.$$

(ii) *Moreover, if there exists $L > 0$ such that for all $t \geq 0$, ℓ_t is L -Lipschitz continuous for $\|\cdot\|$, then the choice*

$$\eta_t = \frac{\eta \sqrt{2K}}{L\sqrt{t+1}}, \quad t \geq 0,$$

for some $\eta > 0$, guarantee for all $T \geq 0$,

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \left(\frac{h(x) - \min h}{\eta} + \eta \right) L \sqrt{\frac{T+1}{2K}}.$$

(iii) *Moreover, if $\sup_{x \in \mathcal{X}} h(x) < +\infty$, then $\eta = \sqrt{\max_{\mathcal{X}} h - \min h}$ yields,*

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq L \sqrt{\frac{2(\max_{\mathcal{X}} h - \min h)(T+1)}{K}}.$$

4.3 Follow the regularized leader

Loss linearization has the advantage of using only a subgradient of each loss function as the input to the algorithm, and not the whole function. On the other hand, this approach forgets about the curvature of the loss functions and convexity inequality (3.6) may be far from tight when the loss functions do have curvature.

The follow the regularized leader (FTRL) algorithm does not rely on such a majorization and instead outputs the regularized minimizer of the past cumulative loss based on the actual loss functions.

Lemma 4.3.1. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper lower semicontinuous convex function and ρ a regularizer on \mathcal{X} such that $\text{dom } f \cap \text{dom } \rho \neq \emptyset$. Then, $f + \rho$ admits a unique minimizer on \mathcal{X} .*

Proof.

□

Definition 4.3.2 (Follow the regularized leader). Let $(\rho_t)_{t \geq 0}$ be a sequence of regularizers on \mathcal{X} and $(\ell_t)_{t \geq 0}$ a sequence of loss functions on \mathcal{X} . Then, the associated *follow the regularized leader* (FTRL) iterates are defined as $x_0 = \arg \min_{x \in \mathcal{X}} \rho_0(x)$ and

$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \left\{ \sum_{s=0}^t \ell_s(x) + \rho_{t+1}(x) \right\}, \quad t \geq 0.$$

Remark 4.3.3. The above iterates are well-defined thanks to Lemma 4.3.1.

Remark 4.3.4. In the case of linear losses, FTRL reduces to DA from Definition 3.2.1.

Proposition 4.3.5 (Regret bound for FTRL). *Let $(x_t)_{t \geq 0}$ be defined as in Definition 4.3.2 and $h_0 = \rho_0 - \min \rho_0$. For all $t \geq 0$, let $g_t \in \partial \ell_t(x_t)$, $y_t = -\sum_{s=0}^{t-1} g_s$ and*

$$h_{t+1/2}(x) = h_{t+1}(x) = \sum_{s=0}^t D_{\ell_s}(x, x_s; g_s) + \rho_{t+1}(x) - \min \rho_{t+1}, \quad x \in \mathbb{R}^d.$$

Then,

- (i) $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ is a sequence of regularizers on \mathcal{X} ,
- (ii) $((x_t, y_t))_{t \geq 0}$ is a sequence of UMD iterates associated with regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ and dual increments $(-g_t)_{t \geq 0}$,
- (iii) for all $T \geq 0$ and $x \in \bigcap_{t \geq 0} \text{dom } \rho_t$,

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \rho_T(x) - \min \rho_T + \sum_{t=0}^T D_t^*,$$

where $D_t^* = h_{t+1}^*(y_{t+1}) - h_t^*(y_t) + \langle g_t, x_t \rangle$.

- (iv) Let $\|\cdot\|$ be a norm in \mathbb{R}^d , $K > 0$, ρ a regularizer on \mathcal{X} which we assume K -strongly convex for $\|\cdot\|$ and $(\eta_t)_{t \geq 0}$ a positive nonincreasing sequence. If $\rho_t = \eta_t^{-1} \rho$ for all $t \geq 0$, then for all $T \geq 0$,

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{\rho(x) - \min_{\mathcal{X}} \rho}{\eta_T} + \frac{1}{2K} \sum_{t=0}^T \eta_t \|g_t\|_*^2.$$

- (v) Moreover, if there exists $L > 0$ such that loss functions $(\ell_t)_{t \geq 0}$ are L -Lipschitz continuous for $\|\cdot\|$, and if $\max_{\mathcal{X}} \rho_T < +\infty$, then the choice

$$\eta_t = \frac{1}{L} \sqrt{\frac{2K (\max_{\mathcal{X}} \rho - \min \rho)}{t+1}}, \quad t \geq 0$$

guarantees

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq L \sqrt{\frac{2(\max_{\mathcal{X}} \rho - \min \rho)(T+1)}{K}}.$$

Proof. (i) $h_0 = \rho_0 - \min \rho_0$ is a regularizer on \mathcal{X} by assumption. For all $t \geq 0$, h_{t+1} is lower semicontinuous as the sum of lower semi-continuous functions. It is strictly convex, as the sum of a strictly convex function (regularizer ρ_{t+1}) and convex functions. Besides, for all $s \geq 0$, because ℓ_s has nonempty subdifferential on \mathcal{X} by assumption, its domain contains \mathcal{X} . Therefore,

$$\text{cl dom } h_{t+1} = \text{cl} \left(\text{dom } \rho_{t+1} \cap \bigcap_{0 \leq s \leq t} \text{dom } \ell_s \right) = \text{cl dom } \rho_{t+1} = \mathcal{X}.$$

It remains to prove that $\text{dom } h_{t+1}^* = \mathbb{R}^d$. For $y \in \mathbb{R}^d$,

$$\begin{aligned} h_{t+1}^*(y) &= \sup_{y \in \mathbb{R}^d} \{ \langle y, x \rangle - h_{t+1}(x) \} \\ &= \sup_{y \in \mathbb{R}^d} \left\{ \langle y, x \rangle - \sum_{s=1}^t D_{\ell_s}(x, x_s; g_s) - \rho_{t+1}(x) + \min \rho_{t+1} \right\} \end{aligned}$$

which is a finite quantity because Lemma 4.3.1 applies. Hence, h_{t+1} is indeed a regularizer on \mathcal{X} .

(ii) For all $t \geq 0$,

$$\begin{aligned} x_t &= \arg \min_{x \in \mathcal{X}} \left\{ \sum_{s=0}^{t-1} \ell_s(x) + \rho_t(x) \right\} \\ &= \arg \min_{x \in \mathcal{X}} \left\{ \sum_{s=0}^{t-1} (\ell_s(x_s) + \langle g_s, x - x_s \rangle + D_{\ell_s}(x, x_s; g_s)) + \rho_t(x) \right\} \\ &= \arg \min_{x \in \mathcal{X}} \left\{ \sum_{s=0}^{t-1} (\langle g_s, x \rangle + D_{\ell_s}(x, x_s; g_s)) + \rho_t(x) \right\} \\ &= \arg \min_{x \in \mathcal{X}} \{ \langle -y_t, x \rangle + h_t(x) \} \\ &= \arg \min_{x \in \mathbb{R}^d} \{ \langle -y_t, x \rangle + h_t(x) \} \\ &= \nabla h_t^*(y_t), \end{aligned}$$

where the penultimate equality holds because for all $t \geq 0$, $\text{dom } h_t \subset \mathcal{X}$. The above is equivalent to $y_t \in \partial h_t^*(x_t)$ by Proposition 1.4.6. Because $h_{t+1/2} = h_{t+1}$, the above also imply for all $t \geq 0$,

$$x_{t+1} = \nabla h_{t+1/2}^*(y_{t+1}) = \nabla h_{t+1/2}^*(y_t - g_t),$$

hence the result.

(iii) Applying Lemma 2.5.2 with notation therein gives for all $t \geq 0$,

$$\langle g_t, x_t - x \rangle \leq D_t - D_{t+1} + D_t^* + (\rho_{t+1}(x) - \min \rho_{t+1}) - (\rho_t(x) - \min \rho_t) + D_{\ell_t}(x, x_t; g_t).$$

Reorganizing the terms gives

$$\ell_t(x_t) - \ell_t(x) = D_t - D_{t+1} + D_t^* + (\rho_{t+1}(x) - \min \rho_{t+1}) - (\rho_t(x) - \min \rho_t).$$

Summing yields

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq D_0 + \sum_{t=0}^T D_t^* + (\rho_{T+1}(x) - \min \rho_{T+1}) - (\rho_0(x) - \min \rho_0).$$

Note that for a given $T \geq 0$, iterates x_0, \dots, x_T do not depend on ρ_{T+1} , and thus, the above analysis is valid with e.g. $\rho_{T+1} = \rho_T$, and we can make the this substitution in the above right-hand side¹. Besides,

$$D_0 = h_0(x) - h_0(x_0) - \langle 0, x - x_0 \rangle = \rho_0(x) - \min \rho_0.$$

Simplifying gives the result.

(iv) Continuing the above analysis (fixing $T \geq 0$ and considering $\rho_{T+1} = \rho_T$, which here corresponds to $\eta_{T+1} = \eta_T$), for all $0 \leq t \leq T$, h_t is K/η_t -strongly convex for $\|\cdot\|$ as the sum of convex functions and a K/η_t -strongly convex function (ρ_t); besides, $h_{t+1} \geq h_t$ because parameters $(\eta_t)_{t \geq 0}$ are nonincreasing. Then,

$$h_{t+1} - h_t = D_{\ell_t}(\cdot, x_t; g_t) + \left(\frac{1}{\eta_{t+1}} - \frac{1}{\eta_t} \right) \rho \geq 0,$$

which implies $h_{t+1}^* \leq h_t^*$. Therefore,

$$\begin{aligned} D_t^* &= h_{t+1}^*(y_t - g_t) - h_t^*(y_t) + \langle g_t, x_t \rangle \\ &\leq h_t^*(y_t - g_t) - h_t^*(y_t) + \langle g_t, x_t \rangle \\ &= D_{h_t^*}(y_t - g_t, y_t) \leq \frac{\eta_t}{2K} \|g_t\|_*^2, \end{aligned}$$

where we used Proposition 1.6.4, hence the result. (v) is then an easy consequence. \square

Remark 4.3.6. The above last regret bound (v) provide a guarantee similar to those given in Section 4.2 by OMD and DA with loss linearization, and thus does not make explicit the advantage of FTRL. However, this bound corresponds to the case where the losses are assumed Lipschitz continuous and therefore may very well have no curvature. In the case where losses do have curvature, above regret bound (iii) may be much smaller, and FTRL performs in practice much better and OMD and DA.

¹This argument remains valid for any choice of ρ_{T+1} , but we should keep in mind that D_T^* depends on ρ_{T+1} .

4.4 Strongly convex losses

We now assume that loss functions have strong convexity and establish below that this can be leveraged to achieve much smaller regret bounds of order $\log T$ instead of \sqrt{T} .

Example 4.4.1 (Square loss). Square loss functions of the form $\ell(x) = \frac{1}{2} \|x - z\|_2^2$ for some $z \in \mathbb{R}^d$ are 1-strongly convex with for $\|\cdot\|_2$ by Corollary 1.6.8.

If the decision maker has access, after having chosen x_t , to a subgradient $g_t \in \partial \ell_t(x_t)$ and to the strong convexity parameter of loss function ℓ_t , it is enough to use online gradient descent with a well-chosen step-size that depends on the strong convexity parameters to obtain the following guarantee.

Proposition 4.4.2 (OGD with strongly convex losses). *Let $(K_t)_{t \geq 0}$ be a positive sequence, $(\ell_t)_{t \geq 0}$ loss functions such that for all $t \geq 0$, ℓ_t is K_t -strongly convex for $\|\cdot\|_2$ and $x_0 \in \mathcal{X}$. Consider online gradient descent iterates:*

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t - \gamma_t g_t), \quad t \geq 0,$$

where $g_t \in \partial \ell_t(x_t)$ and $\gamma_t = (\sum_{s=0}^t K_s)^{-1}$. Then, for all $T \geq 0$,

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{1}{2} \sum_{t=0}^T \frac{\|g_t\|_2^2}{\sum_{s=0}^t K_s}.$$

In particular, if there exists $K, L > 0$ such that for all $t \geq 0$, $K_t = K$ and ℓ_t is L -Lipschitz continuous for $\|\cdot\|_2$, then for all $T \geq 0$,

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{L^2}{2K} (1 + \log(T + 1)).$$

Proof. Using the characterization of strong convexity from Proposition 1.6.4, we write

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \sum_{t=0}^T \left(\langle g_t, x_t - x \rangle - \sum_{t=0}^T \frac{K_t}{2} \|x - x_t\|_2^2 \right).$$

We then conclude by applying Proposition 3.3.13 and simplifying. \square

The above result will be used to derive convergence guarantees for SGD the context of empirical risk minimization with convex losses and Ridge regularization.

We now turn to the *follow the leader* (FTL not FTRL) algorithm which guarantees the same regret bound as OMD, but for any norm $\|\cdot\|$, without requiring the knowledge of the norm $\|\cdot\|$ nor of the strong convexity parameters. This is therefore a much stronger guarantee.

Proposition 4.4.3 (FTL with strongly convex losses). *Let $\|\cdot\|$ be a norm in \mathbb{R}^d , $(K_t)_{t \geq 0}$ a positive sequence, and $(\ell_t)_{t \geq 0}$ loss functions on \mathcal{X} such that for all $t \geq 0$, ℓ_t is K_t -strongly convex for $\|\cdot\|$ and $x_0 \in \mathcal{X}$. Consider*

$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \left\{ \sum_{s=0}^t \ell_s(x) \right\}, \quad t \geq 0. \quad (4.2)$$

Let $h_0 = \varepsilon \|x - x_0\|_2^2 + I_{\mathcal{X}}(x)$ (for some $\varepsilon > 0$) and for all $t \geq 0$,

$$h_{t+1/2}(x) = h_{t+1}(x) = \sum_{s=0}^t D_{\ell_s}(x, x_s; g_s) + I_{\mathcal{X}}(x),$$

where for all $s \geq 0$, $g_s \in \partial \ell_s(x_s)$. Then,

(i) $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ is a sequence of regularizers on \mathcal{X} ,

(ii) $((x_t, -\sum_{s=1}^{t-1} g_s))_{t \geq 0}$ is a sequence of UMD iterates associated with regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ and dual increments $(-g_t)_{t \geq 0}$,

(iii) for all $T \geq 0$,

$$\max_{x \in \mathcal{X}} \sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{1}{2} \sum_{t=0}^T \frac{\|g_t\|_*^2}{\sum_{s=0}^t K_s}.$$

Proof. The FTL iterates are well-defined because for $t \geq 0$, $\sum_{s=0}^t \ell_s + I_{\mathcal{X}}$ is a proper lower semicontinuous strongly convex function, and thus admits a unique minimizer by Proposition 1.6.12.

(i) Let $t \in \frac{1}{2}\mathbb{N}$. h_t is lower semicontinuous as the sum of lower semicontinuous functions. It is also strictly convex because it is strongly convex. For $t = 0$, $\text{dom } h_0 = \mathcal{X}$ as an immediate consequence of the definition and for $t \geq 1$, because the domain of the convex losses contain \mathcal{X} ,

$$\mathcal{X} \subset \text{dom} \left(\sum_{s=0}^{t-1} D_{\ell_s}(\cdot, x_s; g_s) \right),$$

and therefore

$$\text{dom } h_t = \text{dom} \left(\sum_{s=0}^{t-1} D_{\ell_s}(\cdot, x_s; g_s) + I_{\mathcal{X}} \right) = \mathcal{X}.$$

Besides, strong convexity also ensures that $\text{dom } h_t^* = \mathbb{R}^d$ by Proposition 2.2.3, hence the result.

(ii) Similar to the proof of Proposition 4.3.5.

(iii) For $t \geq 0$, applying Lemma 2.5.2 gives, with notation therein,

$$\langle g_t, x_t - x \rangle \leq D_t - D_{t+1} + D_t^* + D_{\ell_t}(x, x_t; y_t).$$

Rearranging gives

$$\ell_t(x_t) - \ell_t(x) \leq D_t - D_{t+1} + D_t^*.$$

Summing gives

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \varepsilon \|x - x_0\|_2^2 + \sum_{t=0}^T D_t^*.$$

The above being true for all $\varepsilon > 0$, the first term in the above right-hand side can be removed. It remains to bound D_t^* from above. Let $t \geq 0$. Note that $x \mapsto D_{\ell_t}(x, x_t; g_t)$ is a nonnegative function which attains its minimum, which is 0, for $x = x_t$. Then, with notation $y_t = -\sum_{s=0}^{t-1} g_s$ we can write

$$\begin{aligned} x_t &= \nabla h_t^*(y_t) = \arg \min_{x \in \mathbb{R}^d} \left\{ -\langle y_t, x \rangle + \sum_{s=0}^{t-1} D_{\ell_s}(x, x_s; g_s) \right\} \\ &= \arg \min_{x \in \mathbb{R}^d} \left\{ -\langle y_t, x \rangle + \sum_{s=0}^t D_{\ell_s}(x, x_s; g_s) \right\} \\ &= \nabla h_{t+1}^*(y_t) = \nabla h_{t+1/2}^*(y_t), \end{aligned}$$

and similarly $h_t^*(y_t) = h_{t+1/2}^*(y_t) = h_{t+1}^*(y_t)$. Besides, because ℓ_t (and therefore $D_{\ell_t}(\cdot, x_t; g_t)$) is K_t -strongly convex for $\|\cdot\|$ by assumption, $h_{t+1} = \sum_{s=0}^t D_{\ell_t}(\cdot, x_t; g_t)$ is $\sum_{s=0}^t K_s$ -strongly convex. Hence,

$$\begin{aligned} D_t^* &= h_{t+1/2}^*(y_t - g_t) - h_t^*(y_t) + \langle g_t, x_t \rangle \\ &= h_{t+1}^*(y_t - g_t) - h_{t+1}^*(y_t) + \langle g_t, \nabla h_{t+1}^*(y_t) \rangle \\ &= D_{h_{t+1}^*}(y_t - g_t, y_t) \\ &\leq \frac{\|g_t\|_*^2}{2 \sum_{s=0}^t K_s}, \end{aligned}$$

where the inequality holds by Corollary 1.6.5, hence the result. \square

4.5 Online linear regression

We consider the following online linear regression problem, which does not exactly fit in the framework of online convex optimization. At each step $t \geq 0$,

- Nature chooses and reveals $w_t \in \mathbb{R}^d$,

- the Decision Maker chooses $x_t \in \mathbb{R}^d$,
- Nature chooses and reveals $z_t \in \mathbb{R}$,
- the Decision Maker incurs loss $\frac{1}{2}(\langle w_t, x_t \rangle - z_t)^2$.

The corresponding regret then writes

$$\sum_{t=0}^T \left(\frac{1}{2}(\langle w_t, x_t \rangle - z_t)^2 - \frac{1}{2}(\langle w_t, x \rangle - z_t)^2 \right), \quad T \geq 0, \quad x \in \mathbb{R}^d.$$

The following algorithm is a variant of FTRL which at step $t \geq 0$ uses the knowledge of w_t to choose x_t . Since the loss function at step $t \geq 0$ writes

$$\ell_t(x) = \frac{1}{2} \langle w_t, x \rangle^2 - z_t \langle w_t, x \rangle + \frac{1}{2} z_t^2,$$

the algorithm chooses x_t by minimizing the past cumulatives loss functions *plus* the known part of the next loss function ℓ_t , meaning the first above term $\frac{1}{2} \langle w_t, x \rangle^2$, *plus* a regularization term $\frac{\lambda}{2} \|x\|_2^2$.

Definition 4.5.1 (Vovk–Azoury–Warmuth algorithm). Let $\lambda > 0$, and $((w_t, z_t))_{t \geq 0}$ a sequence in $\mathbb{R}^d \times \mathbb{R}$. The associated *Vovk–Azoury–Warmuth iterates* are defined as

$$x_t = \arg \min_{x \in \mathbb{R}^d} \left\{ \frac{1}{2} \sum_{s=0}^{t-1} (z_s - \langle w_s, x \rangle)^2 + \frac{1}{2} \langle w_t, x \rangle^2 + \frac{\lambda}{2} \|x\|_2^2 \right\}, \quad t \geq 0.$$

Lemma 4.5.2 (Lemma 1.11 and Theorem 11.7 in [CBL06]). Let $T \geq 0$, $w_0, \dots, w_T \in \mathbb{R}^d$ and $\lambda > 0$. For all $0 \leq t \leq T$, denote $S_t = \lambda I + \sum_{s=0}^t w_s w_s^\top$. Then,

$$\sum_{t=0}^T w_t^\top S_t^{-1} w_t \leq \sum_{i=1}^d \log \left(1 + \frac{\lambda_i}{\lambda} \right),$$

where $\lambda_1, \dots, \lambda_d$ are the eigenvalues of $S_T - \lambda I$.

Proposition 4.5.3. Consider the iterates $(x_t)_{t \geq 0}$ defined as Definition 4.5.1, $h_0(x) = \frac{\lambda}{2} \|x\|_2^2$ and

$$h_{t+1/2}(x) = h_{t+1}(x) = \frac{1}{2} x^\top \left(\lambda I + \sum_{s=0}^{t+1} w_s w_s^\top \right) x, \quad t \geq 0.$$

Then,

- (i) $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ is a sequence of regularizers on \mathbb{R}^d ,

(ii) $((x_t, \sum_{s=0}^{t-1} z_s w_s))_{t \geq 0}$ is a sequence of UMD iterates associated with $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ and dual increments $(z_t w_t)_{t \geq 0}$,

(iii) for all $T \geq 0$,

$$\sum_{t=0}^T (\ell_t(x_t) - \ell_t(x)) \leq \frac{\lambda}{2} \|x\|_2^2 + \frac{dZ_T^2}{2} \log \left(1 + \frac{W_T^2 T}{\lambda d} \right).$$

where

$$Z_T = \max_{0 \leq t \leq T} |z_t| \quad \text{and} \quad W_T = \max_{0 \leq t \leq T} \|w_t\|_2.$$

Proof. (i) For $t \geq 0$, $\frac{\lambda}{2}I + \sum_{s=0}^t w_s w_s^\top$ is symmetric positive definite as the sum of a positive definite matrix ($\frac{\lambda}{2}I$) and positive semi-definite matrices. h_t is therefore the corresponding squared Mahalanobis norm, which is indeed a regularizer on \mathbb{R}^d .

(ii) For $t \geq 0$, denote $y_t = \sum_{s=0}^{t-1} z_s w_s$. Then,

$$\begin{aligned} x_t &= \arg \min_{x' \in \mathbb{R}^d} \left\{ \frac{1}{2} \sum_{s=0}^{t-1} (\langle w_s, x' \rangle - z_s)^2 + \frac{1}{2} \langle w_t, x' \rangle^2 + \frac{\lambda}{2} \|x'\|_2^2 \right\} \\ &= \arg \min_{x' \in \mathbb{R}^d} \left\{ \frac{1}{2} \sum_{s=0}^{t-1} (\langle w_s, x' \rangle^2 - 2z_s \langle w_s, x' \rangle + z_s^2) + \frac{1}{2} \langle w_t, x' \rangle^2 + \frac{\lambda}{2} \|x'\|_2^2 \right\} \\ &= \arg \min_{x' \in \mathbb{R}^d} \left\{ - \sum_{s=0}^{t-1} \langle z_s w_s, x' \rangle + \frac{1}{2} \sum_{s=0}^t \langle w_s, x' \rangle^2 + \frac{\lambda}{2} \|x'\|_2^2 \right\} \\ &= \nabla h_t^*(y_t), \end{aligned}$$

which can also be written $y_t \in \partial h_t(x_t)$. Then, it also holds that

$$x_{t+1} = \nabla h_{t+1}^*(y_{t+1}) = \nabla h_{t+1/2}^*(y_{t+1}) = \nabla h_{t+1/2}^*(y_t + z_t w_t),$$

hence the result.

(iii) Applying Lemma 2.5.2 gives with notation therein

$$\langle z_t w_t, x - x_t \rangle \leq D_t - D_{t+1} + D_t^* + h_{t+1}(x) - h_t(x), \quad t \geq 0.$$

Summing gives

$$\sum_{t=0}^T \langle z_t w_t, x - x_t \rangle \leq D_0 + \sum_{t=0}^T D_t^* + h_{T+1}(x) - h_0(x).$$

Because iterates x_0, \dots, x_T do not depend on h_{T+1} , the above analysis can be carried by h_{T+1} replaced by h_T . Besides, $x_0 = y_0 = 0$ by definition, which implies $D_0 = h_0(x)$. Therefore,

$$\sum_{t=0}^T \langle z_t w_t, x - x_t \rangle \leq \sum_{t=0}^T D_t^* + \frac{\lambda}{2} \|x\|_2^2 + \frac{1}{2} x^\top \left(\sum_{t=0}^T w_t w_t^\top \right) x. \quad (4.3)$$

For each $t \geq 0$, denote $S_t = \frac{\lambda}{2}I + \sum_{s=0}^t w_s w_s^\top$ and using Fenchel's inequality from Remark 1.4.3,

$$\begin{aligned}
D_t^* &= h_{t+1}^*(y_{t+1}) - h_t^*(y_t) - \langle z_t w_t, x_t \rangle \\
&= D_{h_t^*}(y_{t+1}, y_t) + h_{t+1}^*(y_{t+1}) - h_t^*(y_{t+1}) \\
&\leq D_{h_t^*}(y_{t+1}, y_t) + \langle y_{t+1}, x_{t+1} \rangle - h_{t+1}(x_{t+1}) - \langle y_{t+1}, x_{t+1} \rangle + h_t(x_{t+1}) \\
&= \frac{1}{2}(z_t w_t)^\top S_t^{-1}(z_t w_t) - \langle w_{t+1}, x_{t+1} \rangle^2,
\end{aligned} \tag{4.4}$$

where we expressed the Bregman divergence of a squared Mahalanobis norm. Besides, the regret with respect to the actual loss function can be written as follows:

$$\begin{aligned}
&\frac{1}{2} \left((\langle w_t, x_t \rangle - z_t)^2 - (\langle w_t, x \rangle - z_t)^2 \right) \\
&= \frac{1}{2} \langle w_t, x_t \rangle^2 - \frac{1}{2} \langle w_t, x \rangle^2 + z_t \langle w_t, x - x_t \rangle.
\end{aligned} \tag{4.5}$$

Combining (4.3), (4.4) and (4.5) together with Lemma 4.5.2 gives the result. \square

Chapter 5

Blackwell's approachability

Blackwell's approachability is a very general and somewhat abstract framework for sequential decision problems, where the Decision Maker obtains after each step an outcome vector in e.g. \mathbb{R}^d , that depends on its decision and the decision of Nature through an *outcome function*. The original question was the following: given a subset of \mathbb{R}^d called the *target*, and assuming that the outcome function is bounded, what are the conditions on the outcome function and target set such that the Decision Maker has an algorithm which guarantees that the average outcome vector converges to the target set? And when such a conditions is satisfied, what algorithm does guarantee such a convergence? This framework contains regret minimization, many variants of regret minimization, and other problems such as asymptotic calibration in statistics.

In our presentation, we restrict to target sets that are closed convex cones, which contain all special cases that are of interest to us, and simplifies the link with the tools from previous chapters. This link allows in Section 5.4 below the conversion of online linear optimization algorithms into approachability algorithms. Then in Section 5.5, the problem of regret minimization on the simplex is revisited from an approachability point of view and new algorithms are derived. Among them are the important regret matching (RM) and regret matching+ (RM+) algorithms which demonstrate excellent performance in practice, in particular in the context of learning in games. In Section 5.6, several other applications are presented.

We start with a few definitions and properties about closed convex cones.

5.1 Closed convex cones

The proofs are left as exercises.

Definition 5.1.1. A nonempty set $\mathcal{C} \subset \mathbb{R}^d$ is a *closed convex cone* if it is closed and if for all $x, x' \in \mathcal{C}$ and $\lambda \in \mathbb{R}_{\geq 0}$, $x + x' \in \mathcal{C}$ and $\lambda x \in \mathcal{C}$.

Proposition 5.1.2. *A closed convex cone is convex.*

Remark 5.1.3. A closed convex cone \mathcal{C} being nonempty, closed and convex, the Euclidean projection onto \mathcal{C} is well-defined and denoted $\Pi_{\mathcal{C}}$.

Proposition 5.1.4. *For all $x \in \mathbb{R}^d$ and $\lambda \geq 0$, $\Pi_{\mathcal{C}}(\lambda x) = \lambda \Pi_{\mathcal{C}}(x)$.*

Definition 5.1.5. Let $\mathcal{A} \subset \mathbb{R}^d$. The *polar cone* of \mathcal{A} is the set

$$\mathcal{A}^\circ = \left\{ y \in \mathbb{R}^d, \forall x \in \mathcal{A}, \langle y, x \rangle \leq 0 \right\}.$$

Proposition 5.1.6. *A polar cone is a closed convex cone.*

Proposition 5.1.7. *If \mathcal{C} is a closed convex cone, then $\mathcal{C}^{\circ\circ} = \mathcal{C}$.*

Proposition 5.1.8. *The negative orthant \mathbb{R}_-^d is a closed convex cone and its polar cone is the positive orthant \mathbb{R}_+^d .*

Theorem 5.1.9 (Moreau's decomposition theorem). *Let $\mathcal{C} \subset \mathbb{R}^d$ be a closed convex cone. For all $x \in \mathbb{R}^d$, it holds that:*

$$x = \Pi_{\mathcal{C}}(x) + \Pi_{\mathcal{C}^\circ}(x),$$

where $\Pi_{\mathcal{C}}$ and $\Pi_{\mathcal{C}^\circ}$ denote the Euclidean projection onto \mathcal{C} and \mathcal{C}° respectively.

5.2 Framework

Let \mathcal{A}, \mathcal{B} be two nonempty sets with no particular structure, and $g : \mathcal{A} \times \mathcal{B} \rightarrow \mathbb{R}^d$. The elements of \mathcal{A} (resp. \mathcal{B}) are called the *actions* of the Decision Maker (resp. of Nature). g is called the *outcome function*. At step $t \geq 0$,

- the Decision Maker chooses action $a_t \in \mathcal{A}$,
- Nature chooses action $b_t \in \mathcal{B}$,
- outcome vector $r_t := g(a_t, b_t) \in \mathbb{R}^d$ is revealed.

We aim at defining algorithms which guarantee some bound on the distance of the average or cumulative outcome vector to a given closed convex cone $\mathcal{C} \subset \mathbb{R}^d$ called the *target*.

With no assumption, it is not possible for the Decision Maker to ensure the convergence of average outcome vectors to the target set, as the outcome function may very well output vectors which are far away from the target. The following assumption corresponds to a favorable case, which in the case where g is bounded, can be proved to be a characterization of the Decision Maker having an algorithm ensuring convergence to \mathcal{C} .

Definition 5.2.1. A closed convex cone \mathcal{C} satisfies *Blackwell's condition* with respect to outcome function g if there exists $\alpha : \mathcal{C}^\circ \rightarrow \mathcal{A}$ such that

$$\forall x \in \mathcal{C}^\circ, \forall b \in \mathcal{B}, \quad \langle g(\alpha(x), b), x \rangle \leq 0.$$

α is then called an *oracle* associated with \mathcal{C} and g .

Remark 5.2.2. It follows from the above definition that when Blackwell's condition is satisfied, there exists an oracle satisfying

$$x' = \lambda x \quad \text{for some } \lambda > 0 \quad \implies \quad \alpha(x') = \alpha(x). \quad (5.1)$$

Remark 5.2.3 (Geometric interpretation of Blackwell's condition). Blackwell's condition means that for any given hyperplane containing the target, the Decision Maker has an action which forces the next outcome vector to belong the same side of the hyperplane as the target, regardless of Nature's action. An equivalent, and more informal interpretation is the following. Starting from a point $y \in \mathbb{R}^d$, the "direction to \mathcal{C} " is $\Pi_{\mathcal{C}}(y) - y$, which by Moreau's decomposition theorem belongs to $-\mathcal{C}^\circ$. \mathcal{C}° is in fact the set of all possible opposite "directions to \mathcal{C} ". Blackwell's condition is then equivalent to saying that: given a "direction to \mathcal{C} ", the Decision Maker can ensure that the outcome vector will not be in the opposite of that direction (in the sense of a negative dot product).

In some situations, it is easier to establish the following equivalent dual condition, which however does not constructively provide an oracle.

Proposition 5.2.4 (Blackwell's dual condition). *Assume that \mathcal{A} and \mathcal{B} are convex sets of finite dimensional vector spaces such that \mathcal{A} is compact and outcome function g is bi-affine. Then, a closed convex cone $\mathcal{C} \subset \mathbb{R}^d$ satisfies Blackwell's condition with respect to g if, and only if,*

$$\forall b \in \mathcal{B}, \exists a \in \mathcal{A}, \quad g(a, b) \in \mathcal{C}.$$

Proof. Blackwell's condition can be written

$$\max_{x \in \mathcal{C}^\circ} \min_{a \in \mathcal{A}} \max_{b \in \mathcal{B}} \langle g(a, b), x \rangle \leq 0.$$

Since the above dot product is affine in each of the variables a , b and x , by applying Sion's minimax theorem twice, the above is equivalent to

$$\max_{b \in \mathcal{B}} \min_{a \in \mathcal{A}} \max_{x \in \mathcal{C}^\circ} \langle g(a, b), x \rangle \leq 0,$$

which is exactly the dual condition. □

Several examples of problems that fit into this framework are presented in Section 5.6 below.

For the remaining of this chapter, \mathcal{C} will be a closed convex cone satisfying Blackwell's condition with respect to outcome function g and α an associated oracle satisfying condition (5.1).

5.3 Blackwell's algorithm

We consider the framework and notation introduced in the previous section. In the case where the given target satisfies Blackwell's condition and an associated oracle is known, Blackwell's original algorithm is defined as follows.

Definition 5.3.1. The actions given by *Blackwell's algorithm* are defined as

$$a_t = \alpha \left(\Pi_{\mathcal{C}^\circ} \left(\sum_{s=0}^{t-1} r_s \right) \right), \quad t \geq 0,$$

where $\Pi_{\mathcal{C}^\circ}$ denotes the Euclidean projection onto \mathcal{C}° .

Remark 5.3.2 (Geometric interpretation of Blackwell's algorithm). Given the sum of past outcome vectors $R_{t-1} := \sum_{s=0}^{t-1} r_s$, Blackwell's algorithm chooses action a_t that ensures that the next outcome vector r_t will not be in the opposite direction (in the sense of a negative dot product) of R_{t-1} to \mathcal{C} (which corresponds to $\Pi_{\mathcal{C}}(R_{t-1}) - R_{t-1} = -\Pi_{\mathcal{C}^\circ}(R_{t-1})$ by Moreau's decomposition theorem).

Remark 5.3.3. An important feature is that Blackwell's algorithm does not involve any parameter or step-size to be chosen. It is said to be *parameter-free*.

The above statement first gives a general guarantee for Blackwell's algorithm with no additional assumption. In the case where the outcome vectors are bounded, the distance of the cumulative outcome to the target is bounded as \sqrt{T} ; consequently, because of the cone structure of the target, the average outcome vector converges to the target at speed $1/\sqrt{T}$. This result is a special case of the general construction and analysis presented in Section 5.4 below, but we here give an elementary proof.

Proposition 5.3.4 (Guarantees for Blackwell's algorithm). *If $\mathcal{C} \subset \mathbb{R}^d$ is a closed convex cone satisfying Blackwell's condition, Blackwell's algorithm guarantees for all $T \geq 0$,*

$$\min_{r \in \mathcal{C}} \left\| \sum_{t=0}^T r_t - r \right\|_2 \leq \sqrt{\sum_{t=0}^T \|r_t\|_2^2}.$$

Consequently, if there exists $L > 0$ such that $\|r_t\|_2 \leq L$ for all $t \geq 0$,

$$\min_{r \in \mathcal{C}} \left\| \sum_{t=0}^T r_t - r \right\|_2 \leq L\sqrt{T+1}.$$

Proof. For all $t \geq 0$, denote $R_t = \sum_{s=1}^t r_s$ (and $R_{-1} = 0$) and $\Pi_t = \Pi_{\mathcal{C}}(R_t)$.

$$\begin{aligned} \min_{r \in \mathcal{C}} \|R_t - r\|_2^2 &= \|R_t - \Pi_t\|_2^2 \leq \|R_t - \Pi_{t-1}\|_2^2 = \|R_{t-1} + r_t - \Pi_{t-1}\|_2^2 \\ &= \|R_{t-1} - \Pi_{t-1}\|_2^2 + 2 \langle R_{t-1} - \Pi_{t-1}, r_t \rangle + \|r_t\|_2^2. \end{aligned}$$

We bound the above dot product using Moreau's decomposition theorem as follows:

$$\begin{aligned} \langle R_{t-1} - \Pi_{t-1}, r_t \rangle &= \langle R_{t-1} - \Pi_{\mathcal{C}}(R_{t-1}), r_t \rangle \\ &= \langle \Pi_{\mathcal{C}^\circ}(R_{t-1}), g(\alpha(\Pi_{\mathcal{C}^\circ}(R_{t-1})), b_t) \rangle \leq 0, \end{aligned}$$

where the last inequality holds by definition of Blackwell's algorithm and the equality by definition of oracle α . Therefore,

$$\min_{r \in \mathcal{C}} \|R_t - r\|^2 \leq \min_{r \in \mathcal{C}} \|R_{t-1} - r\|^2 + \|r_t\|_2^2.$$

The result follows from summing over $t = 0, \dots, T$. \square

5.4 Regret-based approachability algorithms

We present a scheme which converts an online linear optimization algorithm into an approachability algorithm. In particular, we transpose the guarantees of DA with time-dependent parameters from Section 3.2 and of OMD with time-dependent step-sizes.

The following statement gives an alternative expression of the distance to the target, measured by an arbitrary norm.

Proposition 5.4.1. *Let $\|\cdot\|$ be a norm in \mathbb{R}^d and B the corresponding closed unit ball. Then,*

$$\max_{x \in \mathcal{C}^\circ \cap B} \langle y, x \rangle = \min_{y' \in \mathcal{C}} \|y' - y\|_*, \quad y \in \mathbb{R}^d.$$

Proof. Let $y \in \mathbb{R}^d$. Using the definition of the dual norm and Sion's minimax theorem,

$$\inf_{y' \in \mathcal{C}} \|y' - y\|_* = \inf_{y' \in \mathcal{C}} \sup_{x \in B} \langle y - y', x \rangle = \sup_{x \in B} \inf_{y' \in \mathcal{C}} \{ \langle y, x \rangle - \langle y', x \rangle \}.$$

Suppose x does not belong to \mathcal{C}° . Then, there exists $y'_0 \in \mathcal{C}$ such that $\langle y'_0, x \rangle > 0$. \mathcal{C} being closed by multiplication by \mathbb{R}_+ , the quantity $\langle y', x \rangle$ (with $y' \in \mathcal{C}$) can be made arbitrarily large by selecting $y' = \lambda y'_0$ and letting $\lambda \rightarrow +\infty$, and thus the above infimum is equal to $-\infty$. Therefore, we can restrict the above supremum to $\mathcal{C}^\circ \cap B$. We thus have

$$\inf_{y' \in \mathcal{C}} \|y' - y\|_* = \sup_{x \in \mathcal{C}^\circ \cap B} \left\{ \langle y, x \rangle - \sup_{y' \in \mathcal{C}} \langle y', x \rangle \right\}.$$

The above embedded supremum is zero because for $x \in \mathcal{C}^\circ \cap B$ and $y' \in \mathcal{C}$ we obviously have $\langle y', x \rangle \leq 0$, and 0 is attained with $y' = 0$. Finally,

$$\inf_{y' \in \mathcal{C}} \|y' - y\|_* = \sup_{x \in \mathcal{C}^\circ \cap B} \langle y, x \rangle.$$

□

The following lemma relates a general class of quantities that can measure the distance of the sum of outcomes vectors to the target with a quantity that can be interpreted as a regret in an auxiliary online linear optimization problem.

Lemma 5.4.2. *Let $\mathcal{X}_0 \subset \mathbb{R}^d$ a nonempty closed bounded set, $(x_t)_{t \geq 0}$ a sequence in \mathcal{C}° , $(b_t)_{t \geq 0}$ a sequence in \mathcal{B} , and for all $t \geq 0$, $r_t = g(\alpha(x_t), b_t)$. Then for all $T \geq 0$,*

$$\max_{x \in \mathcal{X}_0} \left\langle \sum_{t=0}^T r_t, x \right\rangle \leq \max_{x \in \mathcal{X}_0} \sum_{t=0}^T \langle r_t, x - x_t \rangle.$$

Proof. For all $t \geq 0$, because α is an oracle,

$$\langle r_t, x_t \rangle = \langle g(\alpha(x_t), b_t), x_t \rangle \leq 0.$$

The result follows. □

Our conversion scheme can be summarized as follows.

- Choose closed sets $\mathcal{X}_0 \subset \mathcal{X} \subset \mathcal{C}^\circ$, where \mathcal{X} is nonempty and convex;
- choose an online linear optimization algorithm on \mathcal{X} and
 - use it with $(r_t)_{t \geq 0}$ as payoff vectors,
 - get output $(x_t)_{t \geq 0}$ in \mathcal{X} ,
 - choose actions $a_t = \alpha(x_t)$ ($t \geq 0$) in the initial approachability problem.

Then, according to Lemma 5.4.2, quantity $\max_{x \in \mathcal{X}_0} \left\langle \sum_{t=0}^T r_t, x \right\rangle$ is bounded from above by the regret bound that is offered by the online linear optimization algorithm.

In particular, any UMD iterates with regularizers on \mathcal{X} can be converted. We here focus on two special cases: DA with time-dependent parameters and OMD with time-dependent step-sizes.

Proposition 5.4.3 (DA for approachability). *Let $K, L > 0$, $\|\cdot\|$ a norm in \mathbb{R}^d , closed sets $\mathcal{X}_0 \subset \mathcal{X} \subset \mathcal{C}^\circ$ where \mathcal{X} is convex, and h a regularizer on \mathcal{X} which is K -strongly convex for $\|\cdot\|$. Assume that*

- for all $t \geq 0$,

$$a_t = \alpha \left(\nabla h^* \left(\eta_t \sum_{s=0}^{t-1} r_s \right) \right) \quad \text{where } \eta_t = \sqrt{\frac{2K(\max_{\mathcal{X}_0} h - \min h)}{L^2(t+1)}},$$

- for all $t \geq 0$, $\|r_t\|_* \leq L$,
- $\max_{\mathcal{X}_0} h < +\infty$.

Then for all $T \geq 0$,

$$\max_{x \in \mathcal{X}_0} \left\langle \sum_{t=0}^T r_t, x \right\rangle \leq L \sqrt{\frac{2(\max_{\mathcal{X}_0} h - \min h)(T+1)}{K}}.$$

Proof. Combine Proposition 3.2.6 and Lemma 5.4.2. \square

Proposition 5.4.4 (Blackwell's algorithm is a special case of DA). *Let $(\eta_t)_{t \geq 0}$ be a positive sequence and $h_2 = \frac{1}{2} \|\cdot\|_2^2 + I_{\mathcal{C}^\circ}$ the Euclidean regularizer on \mathcal{C}° . Then, Blackwell's algorithm coincide with dual averaging associated with regularizer h_2 and parameters $(\eta_t)_{t \geq 0}$.*

Proof. For all $t \geq 0$, using Proposition 2.2.5 on the properties of the Euclidean regularizer, the DA algorithm can be rewritten as

$$\begin{aligned} a_t &= \alpha \left(\nabla h_2^* \left(\eta_t \sum_{s=0}^{t-1} r_s \right) \right) = \alpha \left(\Pi_{\mathcal{C}^\circ} \left(\eta_t \sum_{s=0}^{t-1} r_t \right) \right) \\ &= \alpha \left(\eta_t \cdot \Pi_{\mathcal{C}^\circ} \left(\sum_{s=0}^{t-1} r_t \right) \right) = \alpha \left(\Pi_{\mathcal{C}^\circ} \left(\sum_{s=0}^{t-1} r_t \right) \right), \end{aligned}$$

where we used Proposition 5.1.4 for the third equality and where the last inequality holds because oracle α satisfies condition (5.1) by assumption. This indeed coincides with the definition of Blackwell's algorithm from Definition 5.3.1. \square

Remark 5.4.5. Blackwell's algorithm being parameter-free, the above result shows that in the special case of the Euclidean regularizer on \mathcal{C}° , the actions chosen by dual averaging do not depend on the parameters $(\eta_t)_{t \geq 0}$.

Proposition 5.4.6 (OMD for approachability). *Let $K, L, R > 0$, $\|\cdot\|$ a norm in \mathbb{R}^d , closed sets $\mathcal{X}_0 \subset \mathcal{X} \subset \mathcal{C}^\circ$ where \mathcal{X} is convex, H a mirror map compatible with \mathcal{X} which is K -strongly convex for $\|\cdot\|$, and $x_0 \in \mathcal{X} \cap \text{int dom } H$. Assume that*

- for all $t \geq 0$,

$$\begin{aligned}\gamma_t &= \frac{R\sqrt{K}}{L\sqrt{t+1}} \\ x_{t+1} &= \arg \max_{x \in \mathcal{X}} \{\langle \nabla H(x_t) + \gamma_t r_t, x \rangle - H(x)\} \\ a_{t+1} &= \alpha(x_{t+1}),\end{aligned}$$

- for all $t \geq 0$, $\|r_t\|_* \leq L$,
- for all $t \geq 0$, $\max_{x \in \mathcal{X}_0} D_H(x, x_t) \leq R^2$.

Then for all $T \geq 0$,

$$\max_{x \in \mathcal{X}_0} \left\langle \sum_{t=0}^T r_t, x \right\rangle \leq RL \sqrt{\frac{2(T+1)}{K}}.$$

Proof. Combine Proposition 3.3.14 and Lemma 5.4.2. \square

In the case of the Euclidean mirror map, OMD is called Online Gradient Descent (OGD). We now consider the corresponding approachability algorithm which we call *greedy Blackwell*, and which can be considered as the counterpart in the OMD family of Blackwell's algorithm. We establish below the same guarantee as for Blackwell's algorithm in Proposition 5.3.4.

Definition 5.4.7. The actions of the *greedy Blackwell algorithm* are given by $x_0 = 0$, $a_0 = \alpha(0)$ and for $t \geq 0$ by

$$x_{t+1} = \Pi_{\mathcal{C}^\circ}(x_t + r_t) \quad \text{and} \quad a_{t+1} = \alpha(x_{t+1}).$$

Proposition 5.4.8 (Greedy Blackwell is OGD for approachability). *Let $\gamma > 0$. The greedy Blackwell algorithm coincide with OGD on \mathcal{C}° with initial point $x_0 = 0$ and constant step-size γ .*

Proof. Denote $(x_t)_{t \geq 0}$ the sequence defined as in Definition 5.4.7 and $(x'_t)_{t \geq 0}$ the sequence from Proposition 5.4.6 associated with Euclidean mirror map $H_2 = \frac{1}{2} \|\cdot\|_2^2$ and $\mathcal{X} = \mathcal{C}^\circ$. Let us prove that $x'_t = \gamma x_t$ for all $t \geq 0$. It is true for $t = 0$, as $x'_0 = x_0 = 0$. Then, by induction, for $t \geq 1$,

$$x'_{t+1} = \Pi_{\mathcal{C}^\circ}(x'_t + \gamma r_t) = \Pi_{\mathcal{C}^\circ}(\gamma x_t + \gamma r_t) = \gamma \cdot \Pi_{\mathcal{C}^\circ}(x_t + r_t) = \gamma x_{t+1},$$

where we used Proposition 5.1.4 for the penultimate equality. The result follows from property (5.1). \square

Remark 5.4.9. Like Blackwell's algorithm, the greedy Blackwell algorithm is parameter-free.

Proposition 5.4.10. *The greedy Blackwell algorithm guarantees for all $T \geq 0$,*

$$\min_{r \in \mathcal{C}} \left\| \sum_{t=0}^T r_t - r \right\|_2 \leq \sqrt{\sum_{t=0}^T \|r_t\|_2^2}.$$

Proof. According to Proposition 5.4.8, the greedy Blackwell algorithm corresponds to OGD on \mathcal{C}° with $x_0 = 0$ and any step-size $\gamma > 0$. Then the DA regret bound from Proposition 3.2.5 (applied with constant mirror map $H = H_2/\gamma$) gives, for all $x \in \mathcal{C}^\circ$ and $T \geq 0$, with notation therein,

$$\sum_{t=0}^T \langle r_t, x - x_t \rangle \leq D_0 + \sum_{t=0}^T \tilde{D}_t^*,$$

where

$$D_0 = \frac{1}{2\gamma} \|x\|_2^2 \quad \text{and} \quad \tilde{D}_t^* = D_{H^*}(\nabla H(x_t) + r_t, \nabla H(x_t)) = \frac{\|r_t\|_2^2}{2\gamma},$$

because the Bregman divergence corresponds to the Euclidean distance in the case of the Euclidean mirror map and because $H^* = \frac{1}{2\gamma} \|\cdot\|_2^2$ as seen in Example 1.4.7. The above is true for all $\gamma > 0$. In particular, $\gamma = \left(\sum_{t=0}^T \|r_t\|_2^2\right)^{-1}$ yields

$$\sum_{t=0}^T \langle r_t, x - x_t \rangle \leq \frac{1}{2} \left(\|x\|_2^2 + 1 \right) \sqrt{\sum_{t=0}^T \|r_t\|_2^2}.$$

Using Lemma 5.4.2 and Proposition 5.4.1 (applied with $\mathcal{X}_0 = \mathcal{C}^\circ \cap B_2$ where B_2 is the closed Euclidean unit ball) then gives the result. \square

5.5 Approachability-based regret minimization on the simplex

In this section, we consider the online linear optimization problem on the simplex and rewrite it as an approachability problem. This approach yields new algorithms. In particular, the Blackwell and greedy Blackwell algorithms give in this case the so-called regret matching (RM) and regret matching+ (RM+) algorithms respectively.

In this section, $(a_t)_{t \geq 0}$ denote the actions of the Decision Maker in the simplex Δ_d and should not be confused with $(x_t)_{t \geq 0}$ which can be interpreted as the output of auxiliary online linear optimization algorithms, which belong to \mathbb{R}_+^d , but not necessarily to Δ_d .

Denote $\mathbb{1} = (1, \dots, 1) \in \mathbb{R}^d$.

Proposition 5.5.1 (Hart–Mas-Colell reduction). *Consider the framework and notation from Section 5.2 with action sets $\mathcal{A} = \Delta_d$, $\mathcal{B} = \mathbb{R}^d$, and outcome function $g : (a, u) \mapsto u - \langle u, a \rangle \mathbb{1}$.*

(i) \mathbb{R}_+^d satisfies Blackwell’s condition with respect to g , with

$$\alpha(x) = \begin{cases} x/\|x\|_1 & \text{if } x \neq 0 \\ a_0 & \text{if } x = 0, \end{cases} \quad x \in \mathbb{R}_+^d,$$

where $a_0 \in \Delta_d$, being an associated oracle.

(ii) Let $(x_t)_{t \geq 0}$ and $(u_t)_{t \geq 0}$ sequences in \mathcal{X} and \mathbb{R}^d respectively, let $a_t = \alpha(x_t)$ and $r_t = g(a_t, u_t)$ for all $t \geq 0$. Then, for all $T \geq 0$,

$$\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - a_t \rangle = \max_{x \in \Delta_d} \left\langle \sum_{t=0}^T r_t, x \right\rangle.$$

Proof. (i) Let $x \in (\mathbb{R}_+^d)^\circ = \mathbb{R}_+^d$. If $x = 0$, the dot product from Blackwell’s condition is zero for any value of $\alpha(x)$. If $x \neq 0$, because $\langle \mathbb{1}, x \rangle = \|x\|_1$, for all $u \in \mathbb{R}^d$,

$$\begin{aligned} \langle g(\alpha(x), u), x \rangle &= \left\langle g\left(\frac{x}{\|x\|_1}, u\right), x \right\rangle = \left\langle u - \left\langle u, \frac{x}{\|x\|_1} \right\rangle \mathbb{1}, x \right\rangle \\ &= \langle u, x \rangle - \left\langle u, \frac{x}{\|x\|_1} \right\rangle \langle \mathbb{1}, x \rangle = \langle u, x \rangle - \langle u, x \rangle = 0. \end{aligned}$$

(ii) For all $x \in \Delta_d$,

$$\begin{aligned} \left\langle \sum_{t=0}^T r_t, x \right\rangle &= \left\langle \sum_{t=0}^T g(u_t, a_t), x \right\rangle = \sum_{t=0}^T \langle u_t - \langle u_t, a_t \rangle \mathbb{1}, x \rangle \\ &= \sum_{t=0}^T (\langle u_t, x \rangle - \langle u_t, a_t \rangle \langle \mathbb{1}, x \rangle) = \sum_{t=0}^T \langle u_t, x - a_t \rangle. \end{aligned}$$

□

In the approachability problem described in Proposition 5.5.1, we now examine the Blackwell’s algorithm and the greedy Blackwell algorithm. They are first defined below with their simplest expressions and then related to Definitions 5.3.1 and 5.4.7.

For $y \in \mathbb{R}^d$, denote y_+ the corresponding vector obtained by taking the positive part in each component, in other words,

$$y_+ = \left(\max(y_i, 0) \right)_{1 \leq i \leq d}.$$

Definition 5.5.2. Let $(u_t)_{t \geq 0}$ be a sequence in \mathbb{R}^d and $a_0 \in \Delta_d$.

- The *regret matching* (RM) algorithm gives for all $t \geq 0$,

$$a_t = \begin{cases} \frac{y_{t,+}}{\|y_{t,+}\|_1} & \text{if } y_{t,+} := \left(\sum_{s=0}^{t-1} (u_s - \langle u_s, a_s \rangle \mathbb{1}) \right)_+ \neq 0, \\ a_0 & \text{otherwise.} \end{cases} \quad t \geq 0.$$

- The *regret matching+* (RM+) algorithm is defined as $x_0 = 0$ and for all $t \geq 0$,

$$a_t = \begin{cases} \frac{x_t}{\|x_t\|_1} & \text{if } x_t \neq 0 \\ a_0 & \text{otherwise,} \end{cases}$$

$$x_{t+1} = (x_t + u_t - \langle u_t, a_t \rangle \mathbb{1})_+.$$

Proposition 5.5.3. (i) RM (resp. RM+) corresponds to Blackwell's algorithm (resp. the greedy Blackwell algorithm) in the approachability problem and oracle described in Proposition 5.5.1.

(ii) Against a sequence of payoff vectors $(u_t)_{t \geq 0}$ in \mathbb{R}^d , with notation from Definition 5.5.2, both RM and RM+ guarantee for all $T \geq 0$,

$$\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - a_t \rangle \leq \sqrt{\sum_{t=0}^T \|u_t - \langle u_t, a_t \rangle \mathbb{1}\|_2^2}.$$

(iii) Moreover, if there exists $L > 0$ such that $\|u_t\|_\infty \leq L$ for all $t \geq 0$, both RM and RM+ guarantee for all $T \geq 0$,

$$\max_{x \in \Delta_d} \sum_{t=0}^T \langle u_t, x - a_t \rangle \leq 2L\sqrt{d(T+1)}.$$

Proof. (i) Because $(\mathbb{R}_-^d)^\circ = \mathbb{R}_+^d$, and considering the oracle α from Proposition 5.5.1, establishing $\Pi_{\mathbb{R}_+^d}(y) = y_+$ for all $y \in \mathbb{R}^d$ will make the definition of Blackwell's algorithm (resp. the greedy Blackwell algorithm) coincide with RM (resp. RM+). Let $y \in \mathbb{R}^d$.

$$\begin{aligned} \Pi_{\mathbb{R}_+^d}(y) &= \arg \min_{y' \in \mathbb{R}_+^d} \|y' - y\|_2^2 = \arg \min_{y'_1, \dots, y'_d \geq 0} \sum_{i=1}^d (y'_i - y_i)^2 \\ &= \left(\arg \min_{y_i \geq 0} (y'_i - y_i)^2 \right)_{1 \leq i \leq d} = \left(\max(y_i, 0) \right)_{1 \leq i \leq d} \\ &= y_+. \end{aligned}$$

(ii) then follows by noticing that $\Delta_d \subset \mathbb{R}_+^d \cap B_2$ (where B_2 is the closed unit Euclidean ball), and combining Proposition 5.4.1 with the guarantees for RM and RM+ from Propositions 5.3.4 and 5.4.8.

(iii) simply follows by writing for $t \geq 0$,

$$\begin{aligned} \|u_t - \langle u_t, a_t \rangle \mathbb{1}\|_2 &\leq \|u_t\|_2 + \langle u_t, a_t \rangle \|\mathbb{1}\|_2 \leq \sqrt{d} \|u_t\|_\infty + \|u_t\|_\infty \|a_t\|_1 \|\mathbb{1}\|_2 \\ &\leq \sqrt{d} \|u_t\|_\infty + \|u_t\|_\infty \|a_t\|_1 \sqrt{d} \leq 2L\sqrt{d}. \end{aligned}$$

□

Remark 5.5.4. The regret bound of order \sqrt{dT} from (iii) in the case where the payoff vectors $(u_t)_{t \geq 0}$ are bounded by L with respect to $\|\cdot\|_\infty$ is suboptimal: recall that the exponential weights algorithm achieves the optimal $\sqrt{T \log d}$ regret bound in Proposition 3.4.4. However, the latter bound needs prior knowledge of L for use in the value of the parameters. In contrast, RM and RM+ are parameter-free and achieve the interestingly adaptive regret bound (ii) with no prior knowledge whatsoever.

The RM and RM+ regret minimization algorithm are of high importance because of their simplicity, ease of implementation, their adaptive character, and their excellent practical performance, in particular in the context of learning in games.

5.6 Further applications

We quickly mention a few problems that are variants of regret minimization and that can be solved using regret-based approachability algorithm from Section 5.4 and that either cannot be solved directly using online linear/convex optimization algorithms, or that can but for which the latter do not directly give optimal guarantees. The approach has a systematic character:

- find actions sets \mathcal{A}, \mathcal{B} , outcome function $g : \mathcal{A} \times \mathcal{B} \rightarrow \mathbb{R}^d$, and closed convex cone \mathcal{C} satisfying Blackwell's condition with respect to g ;
- find set $\mathcal{X}_0 \subset \mathcal{C}^\circ$ such that for sequences of actions $(a_t)_{t \geq 0}$ and $(b_t)_{t \geq 0}$ in \mathcal{A} and \mathcal{B} respectively, quantity

$$\max_{x \in \mathcal{X}_0} \left\langle \sum_{t=0}^T g(a_t, b_t), x \right\rangle$$

is equal or is an upper bound on the quantity of interest in the initial problem;

- choose a closed convex set \mathcal{X} such that $\mathcal{X}_0 \subset \mathcal{X} \subset \mathcal{C}^\circ$ and a sequence of regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ on \mathcal{X} ;

- consider UMD iterates associated with regularizers $(h_t)_{t \in \frac{1}{2}\mathbb{N}}$ and dual increments $(g(a_t, b_t))_{t \geq 0}$;
- transpose corresponding guarantee using Lemma 5.4.2.

We only present below each problem and its reduction to an approachability problem. The choice of an algorithm in the latter and the derivation of the corresponding guarantee are left as exercises.

Regret minimization with global costs The problem of regret minimization with global costs is motivated by load balancing and job scheduling, where at each step, the Decision Maker first chooses a distribution (task allocation) over d machines, and then observes the cost of using each machine, which may be different for each machine and each step. The goal of the Decision Maker is to minimize, not the sum of the cumulative costs of using each machine, but a given function of the vector of cumulative costs. A typical example of such *global cost* function is the ℓ_p norm, which includes as special cases the sum of the costs (for $p = 1$), as well as the makespan i.e. the highest cumulative cost (for $p = \infty$).

Assume $d \geq 2$ and let $\|\cdot\|$ be a norm on \mathbb{R}^d . At step $t \geq 0$,

- the Decision Maker chooses $a_t \in \Delta_d$;
- Nature chooses loss vector $\ell_t \in [0, 1]^d$.

The Decision Maker aims at minimizing the following regret:

$$\left\| \sum_{t=0}^T a_t \odot \ell_t \right\| - \min_{a \in \Delta_d} \left\| \frac{1}{T} \sum_{t=0}^T a \odot \ell_t \right\|, \quad T \geq 0,$$

where \odot denotes the component-wise multiplication. At each step $t \geq 0$, the i -th component of vector $a_t \odot \ell$ is equal to $a_{t,i} \ell_i$ which corresponds to the cost of using machine i for a fraction $a_{t,i}$ of the job. The regret is the difference between the actual global cost incurred by the Decision Maker and the best possible global cost in hindsight for a static distribution $a \in \Delta_d$. Important special cases include the makespan which corresponds to $\|\cdot\| = \|\cdot\|_\infty$: the global cost is then the highest average cost over the machines; and for $\|\cdot\| = \|\cdot\|_1$ the global cost simply corresponds to the sum of the costs of all the machines, and the problem then reduces to basic regret minimization on the simplex.

This problem can be reduced to an approachability problem by considering $\mathcal{A} = \Delta_d$, $\mathcal{B} = [0, 1]^d$, outcome function $g : \mathcal{A} \times \mathcal{B} \rightarrow (\mathbb{R}^d)^2$ defined as

$$g(a, \ell) = (a \odot \ell, \ell), \quad a \in \Delta_d, \ell \in [0, 1]^d,$$

and target set

$$\mathcal{C} = \left\{ (y, y') \in (\mathbb{R}_+^d)^2, \|y\| \leq \min_{a \in \Delta_d} \|a \odot y'\| \right\},$$

which can be proved to satisfy Blackwell's condition.

Online combinatorial optimization Let $d, m \geq 1$ be integers. Let $\mathcal{I} = \{1, \dots, d\}$ and P a set which contains subsets of \mathcal{I} of cardinality m . Denote $\Delta(P)$ the unit simplex in \mathbb{R}^P . At step $t \geq 0$,

- the Decision Maker chooses $a_t \in \Delta(P)$;
- Nature chooses and reveals $v_t \in \mathbb{R}^d$;
- the Decision Maker draws $p_t \sim a_t$ and gets payoff $\sum_{i \in p_t} v_{t,i}$.

The quantity to minimize is the following regret:

$$\max_{p \in P} \sum_{t=0}^T \sum_{i \in p} v_{t,i} - \sum_{t=0}^T \sum_{i \in p_t} v_{t,i}.$$

This problem can be seen as a basic regret minimization problem on finite set P , and payoff vectors $(\sum_{i \in p} v_i)_{p \in P}$ which belong to $[-m, m]^P$ as soon as we assume $v \in [-1, 1]^d$. The exponential weights algorithm would then guarantee a regret bound of order $m\sqrt{T \log |P|}$ by Proposition 3.4.4. However, it is possible to take advantage of the structure of the problem and to construct an algorithm which guarantees a significantly better bound, of order $m\sqrt{T \log(d/m)}$, which is known to be optimal. This is possible by reducing the problem to a well-chosen approachability problem.

Let A be the $d \times |P|$ matrix defined by $A = (\mathbb{1}_{\{i \in p\}})_{\substack{i \in \mathcal{I} \\ p \in P}}$, and for each $p \in P$, let $e_p = (\mathbb{1}_{\{i \in p\}})_{i \in \mathcal{I}} \in \mathbb{R}^d$. We consider action sets $\mathcal{A} = \Delta(P)$ and $\mathcal{B} = [-1, 1]^d$, outcome function

$$g(a, v) = v - \frac{\langle v, a \rangle}{m} \mathbf{1} \in \mathbb{R}^d, \quad a \in \Delta(P), v \in [-1, 1]^d,$$

and target set $\mathcal{C} = A(\Delta(P))^\circ$ where $A(\Delta(P))$ denotes the image of the set $\Delta(P)$ via A seen as a linear map from \mathbb{R}^P to \mathbb{R}^d . \mathcal{C} can then be proved to satisfy Blackwell's condition.

Internal and swap regret We here consider the same framework as in Section 3.4, only the quantity to be minimized is different. At step $t \geq 0$,

- the Decision Maker chooses $a_t \in \Delta_d$,
- Nature chooses and reveals $v_t \in \mathbb{R}^d$,

- the Decision Maker draws $i_t \sim a_t$.

Let Φ be a nonempty subset of $\mathcal{I}^{\mathcal{I}}$. The quantity to minimize is the Φ -regret defined as

$$\max_{\varphi \in \Phi} \sum_{t=0}^T v_{t,\varphi(i_t)} - \sum_{t=0}^T v_{t,i_t},$$

and can be interpreted as follows. For a given map $\varphi \in \Phi$, $\sum_{t=0}^T v_{t,\varphi(i_t)}$ is the cumulative payoff that the Decision Maker would have obtained if he had chosen $\varphi(i)$ each time he actually chose i (for all $i \in \mathcal{I}$). The Φ -regret therefore compares the actual cumulative payoff of the Decision Maker with the best such quantity (for $\varphi \in \Phi$) in hindsight. It is possible to construct an algorithm which guarantees on the Φ -regret a bound of order $\sqrt{T \log |\Phi|}$ by reducing this problem to a well-chosen approachability problem.

Consider action sets $\mathcal{A} = \Delta_d$ and $\mathcal{B} = [-1, 1]^d$, outcome function

$$g(a, v) = \left(\sum_{i \in \mathcal{I}} a_i (v_{\varphi(i)} - v_i) \right)_{\varphi \in \Phi}, \quad a \in \Delta_d, v \in \mathbb{R}^d.$$

and target set \mathbb{R}_-^{Φ} , which can then be proved to satisfy Blackwell's condition.

An important special case is when Φ is the set of all transpositions of \mathcal{I} , in other words, the set of maps $\varphi : \mathcal{I} \rightarrow \mathcal{I}$ such that there exists $i \neq j$ in \mathcal{I} such that

$$\varphi(i) = j, \quad \varphi(j) = i, \quad \text{and} \quad \varphi(k) = k \text{ for all } k \notin \{i, j\}.$$

The Φ -regret is then called the *internal regret* and can be written

$$\max_{i, j \in \mathcal{I}} \sum_{t=0}^T \mathbb{1}_{\{i_t=i\}} (v_{t,j} - v_{t,i}).$$

Chapter 6

Gradient methods in optimization

Let $\mathcal{X} \subset \mathbb{R}^d$ be a nonempty closed convex set.

6.1 Lipschitz convex optimization

Let $L > 0$, $\|\cdot\|$ a norm in \mathbb{R}^d , $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a convex function that is L -Lipschitz continuous for $\|\cdot\|$ and that admits a minimizer on \mathcal{X} , meaning there exists $x_* \in \mathcal{X}$ such that

$$f(x_*) = \min_{x \in \mathcal{X}} f(x).$$

The following statement, when used with $x = x_*$, relates the minimization of f on \mathcal{X} using subgradients with a quantity that can be interpreted as a regret in an online linear optimization problem. All algorithms and results below are then transpositions of familiar online linear optimization algorithms together with their guarantees.

Lemma 6.1.1. *Let $(x_t)_{t \geq 0}$ and $(g_t)_{t \geq 0}$ sequences in \mathbb{R}^d such that $g_t \in \partial f(x_t)$. Then for all positive sequence $(\gamma_t)_{t \geq 0}$, $x \in \mathbb{R}^d$ and $T \geq 0$,*

$$\min_{0 \leq t \leq T} f(x_t) - f(x) \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \sum_{t=0}^T \langle \gamma_t g_t, x_t - x \rangle.$$

Proof. For all $t \geq 0$, the definition a subgradient gives

$$f(x_t) - f(x) \leq \langle g_t, x_t - x \rangle.$$

Multiplying by γ_t and summing over $t = 0, \dots, T$ and dividing by $\sum_{t=0}^T \gamma_t$ gives

$$\frac{\sum_{t=0}^T \gamma_t f(x_t)}{\sum_{t=0}^T \gamma_t} - f(x) \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \sum_{t=0}^T \langle \gamma_t g_t, x_t - x \rangle.$$

The result follows. \square

We first consider the following very general extension of projected (sub)gradient descent, which writes $x_{t+1} = \Pi_{\mathcal{X}}(x_t - \gamma_t g_t)$ ($t \geq 0$), and then derive several corollaries.

Proposition 6.1.2 (Strict UMD iterates for Lipschitz convex optimization). *Let $K, R > 0$, h a regularizer on \mathcal{X} such that $x_* \in \text{dom } h$ and which is K -strongly convex for $\|\cdot\|$, and $((x_t, y_t))_{t \geq 0}$ a sequence of strict UMD iterates associated with regularizer h and dual increments $(-\gamma_t g_t)_{t \geq 0}$ where $g_t \in \partial f(x_t)$ for all $t \geq 0$.*

(i) *Let $T \geq 0$. Then,*

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \left(D_h(x_*, x_0; y_0) + \frac{L^2}{2K} \sum_{t=0}^T \gamma_t^2 \right).$$

(ii) *Let $\gamma > 0$. If $\gamma_t = \gamma \sqrt{2K} / (L\sqrt{t+1})$ for all $t \geq 0$, then*

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{L}{\sqrt{2KT}} \left(\frac{D_h(x_*, x_0; y_0)}{\gamma} + \gamma(1 + \log(T+1)) \right).$$

(iii) *If $D_h(x_*, x_0; y_0) \leq R^2$, then $\gamma = R$ yields,*

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{RL}{\sqrt{2KT}} (2 + \log(T+1)).$$

Corollary 6.1.3 (DA for Lipschitz convex optimization). *Consider the assumptions from Proposition 6.1.2. In particular, the same guarantees hold if:*

$$x_t = \nabla h^* \left(y_0 - \sum_{s=0}^{t-1} \gamma_s g_s \right), \quad t \geq 0,$$

In the context of optimizing a unique objective function, online mirror descent is called mirror descent (MD). Proposition 6.1.2 reduces to the following.

Corollary 6.1.4 (MD for Lipschitz convex optimization). *Let $K, R, \gamma > 0$, $\|\cdot\|$ a norm on \mathbb{R}^d , H a mirror map compatible with \mathcal{X} and K -strongly convex for $\|\cdot\|$, $x_0 \in \mathcal{X} \cap \text{int dom } H$ and for all $t \geq 0$,*

$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \{ \langle \nabla H(x_t) - \gamma_t g_t, x \rangle - H(x) \}$$

where $g_t \in \partial f(x_t)$ and $\gamma_t = \gamma \sqrt{2K} / (L\sqrt{t+1})$.

(i) Let $T \geq 0$. Then,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{L}{\sqrt{2KT}} \left(\frac{D_H(x_*, x_0)}{\gamma} + \gamma(1 + \log(T+1)) \right).$$

(ii) If $D_H(x_*, x_0; y_0) \leq R^2$, then $\gamma = R$ yields,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{RL}{\sqrt{2KT}} (2 + \log(T+1)).$$

Corollary 6.1.5 (Projected GD for Lipschitz convex optimization). *Assume that f is L -Lipschitz continuous for $\|\cdot\|_2$. Let $R, \gamma > 0$, $x_0 \in \mathcal{X}$ and for all $t \geq 0$,*

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t - \gamma_t g_t),$$

where $g_t \in \partial f(x_t)$ and $\gamma_t = \gamma\sqrt{2}/(L\sqrt{t+1})$.

(i) Let $T \geq 0$. Then,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{L}{\sqrt{2T}} \left(\frac{\|x_* - x_0\|_2^2}{2\gamma} + \gamma(1 + \log(T+1)) \right).$$

(ii) If $\frac{1}{2}\|x_* - x_0\|_2^2 \leq R^2$, then $\gamma = R$ yields,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{RL}{\sqrt{2T}} (2 + \log(T+1)).$$

The convergence speed obtained in the above statement is $(\log T)/\sqrt{T}$. Although not an issue in practice, it is possible to shave off the $\log T$ factor using DA with time-dependent parameters instead of time-dependent step-sizes.

Proposition 6.1.6 (DA with time-dependent parameters for Lipschitz convex optimization). *Let $K, R, \eta > 0$, h a regularizer on \mathcal{X} such that $x_* \in \text{dom } h$ and which is K -strongly convex for $\|\cdot\|$, $y_0 \in \mathbb{R}^d$ and*

$$x_t = \nabla h^* \left(\eta_t \left(y_0 - \sum_{i=0}^{t-1} g_i \right) \right), \quad t \geq 0,$$

where $g_t \in \partial f(x_t)$ and $\eta_t = \eta\sqrt{2K}/(L\sqrt{t+1})$ for all $t \geq 0$.

(i) Let $T \geq 0$. Then,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \left(\frac{h(x_*) - \min h}{\eta} + \eta \right) \frac{L}{\sqrt{2K(T+1)}}.$$

(ii) If $h(x_*) - \min h \leq R^2$, then $\eta = R$ yields for all $T \geq 0$,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq LR \sqrt{\frac{2}{K(T+1)}}.$$

Proof. Combine the regret bound for DA with time-dependent parameters from Proposition 3.2.6 with Lemma 6.1.1 applied with constant $\gamma_t = 1$ ($t \geq 0$). \square

All above guarantees need prior knowledge of Lipschitz coefficient L because some parameters of step-sizes must be chosen accordingly to obtain the best bound.

6.2 Smooth convex optimization

Let $L > 0$, $\|\cdot\|$ be a norm in \mathbb{R}^d and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex that is L -smooth for $\|\cdot\|$, and that admits a minimizer on \mathcal{X} , meaning there exists $x_* \in \mathcal{X}$ such that

$$f(x_*) = \min_{x \in \mathcal{X}} f(x).$$

We first give the following very general extension of gradient descent with constant step-size and then derive corollaries.

Proposition 6.2.1. *Let $K > 0$, h a regularizer on \mathcal{X} such that $x_* \in \text{dom } h$ and which is K -strongly convex for $\|\cdot\|$ and $((x_t, y_t))_{t \geq 0}$ a sequence of strict UMD iterates associated with regularizer h and dual increments $(-\frac{K}{L} \nabla f(x_t))_{t \geq 0}$. Then for all $T \geq 0$,*

$$f(x_T) - f(x_*) \leq \frac{L}{K} \frac{D_h(x_*, x_0; y_0)}{T}.$$

Proof. Let $t \geq 0$. The characterization of smoothness from Proposition 1.6.4 gives

$$D_f(x_{t+1}, x_t) \leq \frac{L}{2} \|x_{t+1} - x_t\|^2.$$

Moreover, using the strong convexity of h , we can write

$$D_f(x_{t+1}, x_t) \leq \frac{L}{K} D'_t, \tag{6.1}$$

where $D'_t = D_h(x_{t+1}, x_t; y_t)$. Applying Lemma 2.4.1 for any $x \in \text{dom } h$ gives, with notation therein:

$$D_{t+1} \leq D_t + \frac{K}{L} \langle \nabla f(x_t), x - x_{t+1} \rangle - D'_t.$$

Using (6.1) and the convexity of f , we obtain

$$\begin{aligned} D_{t+1} &\leq D_t + \frac{K}{L} \langle \nabla f(x_t), x - x_t \rangle + \frac{K}{L} \langle \nabla f(x_t), x_t - x_{t+1} \rangle - \frac{K}{L} D_f(x_{t+1}, x_t) \\ &= D_t + \frac{K}{L} (f(x) - f(x_t) - \langle \nabla f(x_t), x_t - x_{t+1} \rangle - D_f(x_{t+1}, x_t)) \\ &= D_t + \frac{K}{L} (f(x) - f(x_{t+1})). \end{aligned}$$

Applying the above for $x = x_t$ (which is possible because $x_t = \nabla h^*(y_t)$ does belong to $\text{dom } h$) gives

$$\begin{aligned} f(x_{t+1}) - f(x_t) &\leq L (D_h(x_t, x_t; y_t) - D_h(x_t, x_{t+1}; y_{t+1})) \\ &= -L \cdot D_h(x_t, x_{t+1}; y_{t+1}) \leq 0. \end{aligned}$$

Then, considering $x = x_*$ (which belongs to $\text{dom } h$ by assumption) and summing gives

$$T (f(x_T) - f(x_*)) \leq \sum_{t=0}^{T-1} (f(x_{t+1}) - f(x_*)) \leq \frac{L}{K} D_h(x_*, x_0; y_0),$$

hence the result. \square

Corollary 6.2.2 (DA for smooth convex optimization). *Let h satisfying the assumptions from Proposition 6.2.1, $y_0 \in \mathbb{R}^d$ and*

$$x_t = \nabla h^* \left(y_0 - \frac{L}{K} \sum_{s=0}^{t-1} \nabla f(x_s) \right), \quad t \geq 0.$$

Then for all $T \geq 0$,

$$f(x_T) - f(x_*) \leq \frac{L}{K} \frac{D_h(x_*, x_0; y_0)}{T}.$$

Corollary 6.2.3 (MD for smooth convex optimization). *Let H be a mirror map compatible with \mathcal{X} and K -strongly convex for $\|\cdot\|$, $x_0 \in \mathcal{X} \cap \text{int dom } H$ and*

$$x_{t+1} = \arg \max_{x \in \mathcal{X}} \left\{ \left\langle \nabla H(x_t) - \frac{K}{L} \nabla f(x_t), x \right\rangle - H(x) \right\}, \quad t \geq 0,$$

Then for all $T \geq 0$,

$$f(x_T) - f(x_*) \leq \frac{L}{K} \frac{D_H(x_*, x_0)}{T}.$$

Corollary 6.2.4 (Projected GD for smooth convex optimization). *Assume that f is L -smooth for $\|\cdot\|_2$. Let $x_0 \in \mathcal{X}$ and*

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t - \frac{K}{L} \nabla f(x_t) \right), \quad t \geq 0.$$

Then for all $T \geq 0$,

$$f(x_T) - f(x_*) \leq \frac{L \|x_0 - x_*\|_2^2}{2T}, \quad T \geq 0.$$

All above guarantees need prior knowledge of L .

6.3 Nesterov's acceleration

We consider the same smooth convex optimization problem as in Section 6.2, where smoothness is with respect to a given norm $\|\cdot\|$. We now define and analyze an extension of Nesterov's accelerated gradient. This algorithm improves the convergence rate from $1/T$ to $1/T^2$.

Proposition 6.3.1 (Accelerated UMD for smooth convex optimization). *Let $K > 0$, h a regularizer on \mathcal{X} such that $x_* \in \text{dom } h$ and which is K -strongly convex for $\|\cdot\|$, $(\gamma_t)_{t \geq 0}$ and $(\lambda_t)_{t \geq 0}$ positive sequences. Let $((v_t, w_t, x_t, w_t))_{t \geq 0}$ a sequence in $(\mathbb{R}^d)^4$ such that*

- $v_0 = w_0 = x_0 = \nabla h^*(y_0)$,
- $((x_t, y_t))_{t \geq 0}$ is a sequence of strict UMD iterates associated with regularizer h and dual increments $(-\gamma_t \nabla f(w_t))_{t \geq 0}$,
- For all $t \geq 0$,

$$w_t = (1 - \lambda_t)v_t + \lambda_t x_t, \tag{6.2}$$

$$v_{t+1} = (1 - \lambda_t)v_t + \lambda_t x_{t+1}. \tag{6.3}$$

where $\gamma_0 = 1/L$ and for all $t \geq 0$,

$$\gamma_{t+1} = \frac{1 + \sqrt{1 + (2L\gamma_t)^2}}{2L}, \quad \text{and} \quad \lambda_t = \frac{1}{L\gamma_t}.$$

Then, for all $T \geq 0$,

$$f(v_T) - f(x_*) \leq \frac{4LD_h(x_*, x_0; y_0)}{KT^2}.$$

With first gather a few properties that are immediate from the above definition.

Lemma 6.3.2. *With assumptions from Proposition 6.3.1, for $t \geq 0$,*

(i) $\lambda_t \in (0, 1)$,

(ii) $w_t - x_t = (\lambda_t^{-1} - 1)(v_t - w_t)$,

(iii) $\gamma_t \lambda_t^{-1} - \gamma_{t+1}(\lambda_{t+1}^{-1} - 1) = 0$,

(iv) For $T \geq 0$, $\gamma_T \lambda_T^{-1} = \sum_{t=0}^T \gamma_t \geq \frac{K(T+1)^2}{4L}$.

Sequence $(x_t, y_t)_{t \geq 0}$ being strict UMD iterates, the following statement follows from Lemma 2.4.1.

Lemma 6.3.3. *With assumptions from Proposition 6.3.1, for $T \geq 0$,*

$$\sum_{t=0}^T \gamma_t \langle \nabla f(w_t), x_{t+1} - x_* \rangle \leq D_h(x_*, x_0; y_0) - \sum_{t=0}^T D_h(x_{t+1}, x_t; y_t).$$

Proof of Proposition 6.3.1. Let $t \geq 0$. Using the definition of smoothness of f between points w_t and v_{t+1} :

$$\begin{aligned} f(v_{t+1}) - f(w_t) &\leq \langle \nabla f(w_t), v_{t+1} - w_t \rangle + \frac{L}{2} \|v_{t+1} - w_t\|^2 \\ &= \lambda_t \langle \nabla f(w_t), x_{t+1} - x_t \rangle + \frac{L\lambda_t^2}{2} \|x_{t+1} - x_t\|^2 \\ &\leq \lambda_t \langle \nabla f(w_t), x_{t+1} - x_t \rangle + \frac{L\lambda_t^2}{K} D_h(x_{t+1}, x_t; y_t), \end{aligned}$$

where we used relation (6.3) from the definition of the algorithm to get the second line, and the K -strong convexity of h (Proposition 1.6.4) to get the third line. Multiplying by $L\lambda_t^2/K$ and simplifying gives:

$$\gamma_t \lambda_t^{-1} (f(v_{t+1}) - f(w_t)) \leq \gamma_t \langle \nabla f(w_t), x_{t+1} - x_t \rangle + D_h(x_{t+1}, x_t; y_t). \quad (6.4)$$

Besides, we can write

$$\begin{aligned} x_{t+1} - x_t &= (x_{t+1} - x_*) + (x_* - w_t) + (w_t - x_t) \\ &= (x_{t+1} - x_*) + (x_* - w_t) + (\lambda_t^{-1} - 1)(v_t - w_t), \end{aligned}$$

where the second line uses relation (ii) from Lemma 6.3.2. Injecting the above into $\langle \nabla f(w_t), x_{t+1} - x_t \rangle$ gives:

$$\begin{aligned} \langle \nabla f(w_t), x_{t+1} - x_t \rangle &= \langle \nabla f(w_t), x_{t+1} - x_* \rangle + \langle \nabla f(w_t), x_* - w_t \rangle \\ &\quad + (\lambda_t^{-1} - 1) \langle \nabla f(w_t), v_t - w_t \rangle \\ &\leq \langle \nabla f(w_t), x_{t+1} - x_* \rangle + f(x_*) - f(w_t) \\ &\quad + (\lambda_t^{-1} - 1) (f(v_t) - f(w_t)). \end{aligned} \quad (6.5)$$

Combining inequalities (7.4) and (7.3), and summing over $t = 0, \dots, T$ gives:

$$\begin{aligned} \sum_{t=0}^T \gamma_t \lambda_t^{-1} (f(v_{t+1}) - f(w_t)) &\leq D_h(x_*, x_0; y_0) - \sum_{t=0}^T D_h(x_{t+1}, x_t; y_t) \\ &+ \sum_{t=0}^T \gamma_t (f(x_*) - f(w_t)) + \sum_{t=0}^T \gamma_t (\lambda_t^{-1} - 1) (f(v_t) - f(w_t)) \\ &+ \sum_{t=0}^T D_h(x_{t+1}, x_t; y_t), \end{aligned}$$

where we used Lemma 6.3.3 to get the first two terms of the right-hand side. Then, simplifying and moving all values of f (except for $f(x_*)$) to the left-hand side, we get:

$$\begin{aligned} \sum_{t=0}^T (\gamma_t + \gamma_t (\lambda_t^{-1} - 1) - \gamma_t \lambda_t^{-1}) f(w_t) &+ \sum_{t=1}^T (\gamma_{t-1} \lambda_{t-1}^{-1} - \gamma_t (\lambda_t^{-1} - 1)) f(v_t) \\ &+ \gamma_0 (\lambda_0^{-1} - 1) f(v_0) + \gamma_T \lambda_T^{-1} f(v_{T+1}) \leq D_h(x_*, x_0; y_0) + \left(\sum_{t=0}^T \gamma_t \right) f(x_*). \end{aligned}$$

The factor in front of $f(w_t)$ is clearly zero, as well as $\gamma_0 (\lambda_0^{-1} - 1)$. The result then follows by applying properties (iii) and (iv) from Lemma 6.3.2. \square

Corollary 6.3.4 (Accelerated dual averaging for smooth convex optimization). *Let $y_0 \in \mathbb{R}^d$, $x_0 = \nabla h^*(y_0)$ and for all $t \geq 0$,*

$$\begin{aligned} w_t &= (1 - \lambda_t)v_t + \lambda_t x_t, \\ x_{t+1} &= \nabla h^* \left(y_0 - \sum_{s=0}^t \gamma_s \nabla f(w_s) \right), \\ v_{t+1} &= (1 - \lambda_t)v_t + \lambda_t x_{t+1}, \end{aligned}$$

where h , $(\gamma_t)_{t \geq 0}$ and $(\lambda_t)_{t \geq 0}$ satisfy the assumptions from Proposition 6.3.1. Then for all $T \geq 0$,

$$f(v_T) - f(x_*) \leq \frac{4L \cdot D_h(x_*, x_0; y_0)}{KT^2}.$$

Proof. Corresponds to Proposition 6.1.2 where $y_{t+1} = y_t - \gamma_t \nabla f(w_t)$ for all $t \geq 0$. \square

Corollary 6.3.5 (Accelerated mirror descent smooth convex optimization). *Let H be a mirror map compatible with \mathcal{X} , K -strongly convex for $\|\cdot\|$, and*

such that $x_* \in \mathcal{X} \cap \text{dom } H$. Let $x_0 \in \mathcal{X} \cap \text{int dom } H$ and for all $t \geq 0$,

$$\begin{aligned} w_t &= (1 - \lambda_t)v_t + \lambda_t x_t, \\ x_{t+1} &= \arg \min_{x \in \mathcal{X}} \{ \langle \nabla H(x_t) - \gamma_t \nabla f(w_t), x \rangle + H(x) \}, \\ v_{t+1} &= (1 - \lambda_t)v_t + \lambda_t x_{t+1}, \end{aligned}$$

where $(\gamma_t)_{t \geq 0}$ and $(\lambda_t)_{t \geq 0}$ satisfy the assumptions from Proposition 6.3.1. Then for all $T \geq 0$,

$$f(v_T) - f(x_*) \leq \frac{4L \cdot D_H(x_*, x_0)}{KT^2}.$$

Proof. Corresponds to Proposition 6.1.2 where $y_t = \nabla H(x_t)$ for all $t \geq 0$. \square

Corollary 6.3.6 (Accelerated gradient descent smooth convex optimization). *Assume that f is L -smooth for $\|\cdot\|_2$. Let $x_0 = v_0 = w_0 \in \mathbb{R}^d$ and for all $t \geq 0$,*

$$\begin{aligned} w_t &= (1 - \lambda_t)v_t + \lambda_t x_t, \\ x_{t+1} &= x_t - \gamma_t \nabla f(w_t), \\ v_{t+1} &= (1 - \lambda_t)v_t + \lambda_t x_{t+1}, \end{aligned}$$

where $(\gamma_t)_{t \geq 0}$ and $(\lambda_t)_{t \geq 0}$ satisfy the assumptions from Proposition 6.3.1. Then, the above can be rewritten as

$$\begin{aligned} v_{t+1} &= w_t - \lambda_t \gamma_t \nabla f(w_t) \\ w_{t+1} &= v_{t+1} + \frac{\lambda_{t+1}(1 - \lambda_t)}{\lambda_t} (v_{t+1} - v_t), \end{aligned}$$

and for all $T \geq 0$,

$$f(v_T) - f(x_*) \leq \frac{2L \|x_0 - x_*\|_2^2}{T^2}.$$

Proof. The definition and the guarantee correspond to Corollary 6.1.4 with H chosen as the Euclidean mirror map and $K = 1$. Besides, for $t \geq 0$,

$$v_{t+1} = w_t + \lambda_t(x_{t+1} - x_t) = w_t - \lambda_t \gamma_t \nabla f(w_t),$$

and

$$\begin{aligned} w_{t+1} &= (1 - \lambda_{t+1})v_{t+1} + \lambda_{t+1}x_{t+1} = v_{t+1} + \lambda_{t+1}(x_{t+1} - v_{t+1}) \\ &= v_{t+1} + \lambda_{t+1}(1 - \lambda_t)(x_{t+1} - v_t) \\ &= v_{t+1} + \frac{\lambda_{t+1}(1 - \lambda_t)}{\lambda_t}(v_{t+1} - v_t). \end{aligned}$$

\square

6.4 Stochastic nonsmooth convex optimization

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a convex function that admits a minimizer on \mathcal{X} , meaning there exists $x_* \in \mathcal{X}$ such that

$$f(x_*) = \min_{x \in \mathcal{X}} f(x).$$

We adapt the approach from Section 6.1 to the case where the algorithm only accesses unbiased stochastic estimators of the (sub)gradients. We present a UMD-based generalization of the celebrated stochastic gradient descent (SGD).

Example 6.4.1 (Finite-sum optimization). Consider an objective function f that is given as

$$f(x) = \frac{1}{n} \sum_{i=1}^n f_i(x),$$

where each function $f_i : \mathbb{R}^d \rightarrow \mathbb{R}$ ($1 \leq i \leq n$) is convex. Important such problems include empirical risk minimization, log-likelihood maximization, etc. A possible way of reducing the per-iteration computational cost is to replace, at each step $t \geq 0$, the computation of an exact (sub)gradient of f at x_t by the computation of a unbiased stochastic estimator by drawing an index i_t uniformly, and considering a subgradient $g_t \in \partial f_{i_t}(x_t)$. Then, it can be verified that indeed $\mathbb{E}[g_t | x_t] \in \partial f(x_t)$.

Lemma 6.4.2. *Let $(x_t)_{t \geq 0}$ and $(g_t)_{t \geq 0}$ be random sequences in \mathbb{R}^d such that for all $t \geq 0$, $\mathbb{E}[g_t | x_t] \in \partial f(x_t)$, and $(\gamma_t)_{t \geq 0}$ a positive sequence, then for all $x \in \mathbb{R}^d$ and $T \geq 0$,*

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x) \right] \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \mathbb{E} \left[\sum_{t=0}^T \langle \gamma_t g_t, x_t - x \rangle \right].$$

Proof. For $t \geq 0$,

$$\begin{aligned} \mathbb{E} \left[\sum_{t=0}^T \langle \gamma_t g_t, x_t - x \rangle \right] &= \mathbb{E} \left[\sum_{t=0}^T \mathbb{E} [\langle \gamma_t g_t, x_t - x \rangle | x_t] \right] \\ &= \mathbb{E} \left[\sum_{t=0}^T \gamma_t \langle \mathbb{E}[g_t | x_t], x_t - x \rangle \right] \\ &\geq \mathbb{E} \left[\sum_{t=0}^T \gamma_t (f(x_t) - f(x)) \right] \\ &\geq \mathbb{E} \left[\left(\sum_{t=0}^T \gamma_t \right) \left(\min_{0 \leq t \leq T} f(x_t) - f(x) \right) \right], \end{aligned}$$

where the penultimate inequality uses the definition of a subgradient. Hence the result. \square

Proposition 6.4.3 (Stochastic UMD for stochastic nonsmooth convex optimization). *Let $K, R, \sigma > 0$, h a regularizer on \mathcal{X} such that $x_* \in \text{dom } h$ and which is K -strongly convex for $\|\cdot\|$, and random sequences $(x_t)_{t \geq 0}$, $(y_t)_{t \geq 0}$ and $(g_t)_{t \geq 0}$ in \mathbb{R}^d and $(\gamma_t)_{t \geq 0}$ a positive sequence such that*

- (x_0, y_0) is deterministic,
- $((x_t, y_t))_{t \geq 0}$ is almost-surely a sequence of strict UMD iterates associated with regularizer h and dual increments $(-\gamma_t g_t)_{t \geq 0}$,
- $\mathbb{E}[g_t | x_t] \in \partial f(x_t)$ for all $t \geq 0$,
- $\mathbb{E}[\|g_t\|^2 | x_t] \leq \sigma^2$ for all $t \geq 0$.

(i) Let $T \geq 0$. Then,

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \left(D_h(x_*, x_0; y_0) + \frac{\sigma^2}{2K} \sum_{t=0}^T \gamma_t^2 \right).$$

(ii) Let $\gamma > 0$. If $\gamma_t = \gamma \sqrt{2K}/(\sigma \sqrt{t+1})$ for all $t \geq 0$, then

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq \frac{\sigma}{\sqrt{2KT}} \left(\frac{D_h(x_*, x_0; y_0)}{\gamma} + \gamma(1 + \log(T+1)) \right).$$

(iii) If $D_h(x_*, x_0; y_0) \leq R^2$, then $\gamma = R$ yields,

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq \frac{R\sigma}{\sqrt{2KT}} (2 + \log(T+1)).$$

Proof. Lemma 6.4.2 gives

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \mathbb{E} \left[\sum_{t=0}^T \langle \gamma_t g_t, x_t - x_* \rangle \right].$$

Because $x_* \in \text{dom } h$, UMD Lemma (Lemma 2.4.1) bounds the expectation from the above right-hand side as

$$\begin{aligned} \mathbb{E} \left[\sum_{t=0}^T \langle \gamma_t g_t, x_t - x_* \rangle \right] &\leq \mathbb{E} \left[D_h(x_*, x_0; y_0) + \frac{1}{2K} \sum_{t=0}^T \|\gamma_t g_t\|^2 \right] \\ &= D_h(x_*, x_0; y_0) + \frac{1}{2K} \mathbb{E} \left[\sum_{t=0}^T \gamma_t^2 \mathbb{E}[\|g_t\|^2 | x_t] \right] \\ &\leq D_h(x_*, x_0; y_0) + \frac{\sigma^2}{2K} \sum_{t=0}^T \gamma_t^2, \end{aligned}$$

which proves (i). (ii) and (iii) follow. \square

Chapter 7

AdaGrad

We present in this chapter two instances of the AdaGrad family of algorithms. It is one of the most important innovations in the topic of regret minimization because of the adaptive property of its regret bounds and because it has led to great success in practice in optimization and deep learning through its many variants, such as RMSprop, Adam, etc.

Let \mathcal{X} be a nonempty closed convex subset of \mathbb{R}^d .

7.1 Definitions

Let $(u_t)_{t \geq 0}$ be a sequence in \mathbb{R}^d , $x_0 \in \mathcal{X}$ and $\gamma > 0$.

Definition 7.1.1. The associated sequence of *AdaGrad-Norm* iterates on \mathcal{X} is given by

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t + \frac{\gamma}{\sum_{s=0}^t \|u_s\|_2^2} u_t \right), \quad t \geq 0,$$

with convention $0/0 = 0$.

Definition 7.1.2. Let $\varepsilon > 0$. The associated sequence $(x_t)_{t \geq 0}$ of *AdaGrad-Diagonal* iterates on \mathcal{X} is defined for each $t \geq 0$ as

$$x'_{t+1} = \left(x'_{t,i} + \frac{\gamma}{\varepsilon + \sqrt{\sum_{s=0}^t u_{s,i}^2}} u_{t,i} \right)_{1 \leq i \leq d},$$
$$x_{t+1} = \arg \min_{x \in \mathcal{X}} \|x - x'_{t+1}\|_{A_t},$$

where

$$A_t = \text{diag} \left(\varepsilon + \sqrt{\sum_{s=0}^t u_{s,i}^2} \right)_{1 \leq i \leq d}.$$

7.2 Regret bounds

Lemma 7.2.1. *Let $(a_t)_{t \geq 0}$ be a nonnegative sequence. Then for all $T \geq 0$,*

$$\sum_{t=0}^T \frac{a_t}{\sqrt{\sum_{s=0}^t a_s}} \leq 2 \sqrt{\sum_{t=0}^T a_t}.$$

with convention $0/0 = 0$.

Proof. For all $t \geq 0$, denote $b_t = \sum_{s=0}^t a_s$. If $b_t = 0$ for some $t \geq 0$, then $b_s = 0$ for all $s \leq t$ and the corresponding terms in the sum are zero because $(a_t)_{t \geq 0}$ is nonnegative by assumption. Without loss of generality, we assume $a_0 = b_0 > 0$. Then,

$$\begin{aligned} \sum_{t=0}^T \frac{b_t - b_{t-1}}{\sqrt{b_t}} &\leq \sum_{t=0}^T \frac{b_t}{\sqrt{b_t}} - \sum_{t=1}^T \frac{b_{t-1}}{\sqrt{b_t}} = \sum_{t=0}^T \frac{b_t}{\sqrt{b_t}} - \sum_{t=0}^{T-1} \frac{b_t}{\sqrt{b_{t+1}}} \\ &= \sqrt{b_T} + \sum_{t=0}^{T-1} b_t \left(\frac{1}{\sqrt{b_t}} - \frac{1}{\sqrt{b_{t+1}}} \right) \\ &= \sqrt{b_T} + \sum_{t=0}^{T-1} \frac{b_t (\sqrt{b_{t+1}} - \sqrt{b_t})}{\sqrt{b_t} \sqrt{b_{t+1}}} \\ &\leq \sqrt{b_T} + \sum_{t=0}^{T-1} (\sqrt{b_{t+1}} - \sqrt{b_t}) \leq 2\sqrt{b_T}, \end{aligned}$$

hence the result. \square

Lemma 7.2.2. *Let $w_1, \dots, w_d \geq 0$. Then,*

$$\sum_{i=1}^d \sqrt{w_i} = \inf_{\substack{v \in (\mathbb{R}_+^*)^d \\ \|v\|_1 \leq 1}} \sqrt{\sum_{i=1}^d \frac{w_i}{v_i}}.$$

Proof. Let $v_1, \dots, v_d > 0$ such that $\sum_{i=1}^d v_i \leq 1$. Then using Jensen's inequality for $z \mapsto z^2$,

$$\left(\sum_{i=1}^d \left(\frac{v_i}{\sum_{j=1}^d v_j} \right) \left(\frac{\sqrt{w_i}}{v_i} \right) \right)^2 \leq \sum_{i=1}^d \frac{v_i}{\sum_{j=1}^d v_j} \frac{w_i}{v_i^2} \leq \sum_{i=1}^d \frac{w_i}{v_i}.$$

Taking the square root gives

$$\sum_{i=1}^d \sqrt{w_i} \leq \sqrt{\sum_{i=1}^d \frac{w_i}{v_i}}.$$

The above inequality is an equality if $w_1 = \dots = w_d = 0$. Otherwise, let $\varepsilon > 0$ and consider

$$v_i = \frac{\sqrt{w_i + \varepsilon}}{\sum_{j=1}^d \sqrt{w_j + \varepsilon}}, \quad 1 \leq i \leq d,$$

which are positive and satisfy $v_1 + \dots + v_d = 1$. Then,

$$\sqrt{\sum_{i=1}^d \frac{w_i}{v_i}} = \sqrt{\sum_{i=1}^d \frac{w_i}{\sqrt{\varepsilon + w_i}} \sum_{j=1}^d \sqrt{\varepsilon + w_j}},$$

which converges to $\sum_{i=1}^d \sqrt{w_i}$ as $\varepsilon \rightarrow 0^+$, hence the result. \square

Proposition 7.2.3 (Regret bound for AdaGrad-Norm). *Let $(u_t)_{t \geq 0}$ be a sequence in \mathbb{R}^d , $x_0 \in \mathcal{X}$, $\varepsilon, \gamma > 0$ and $(x_t)_{t \geq 0}$ the associated sequence of AdaGrad-Norm iterates on \mathcal{X} .*

(i) *Let $x \in \mathcal{X}$ and $T \geq 0$.*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} + \gamma \right) \sqrt{\sum_{t=0}^T \|u_t\|_2^2}.$$

(ii) *If $R \geq \max_{0 \leq t \leq T} \|x_t - x\|_2$ then $\gamma = R/\sqrt{2}$ yields*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq R \sqrt{2 \sum_{t=0}^T \|u_t\|_2^2}.$$

Proof. If $u_t = 0$ for all $t \geq 0$, the result holds. Otherwise, consider

$$\tau = \min \{t \geq 0, u_t \neq 0\}.$$

Let $(\gamma_t)_{t \geq 0}$ be a positive and nonincreasing sequence defined as

$$\gamma_t = \begin{cases} \frac{\gamma}{\|u_\tau\|_2} & \text{if } t \leq \tau \\ \frac{\gamma}{\sqrt{\sum_{s=0}^t \|u_s\|_2^2}} & \text{if } t \geq \tau. \end{cases}$$

Then, $(x_t)_{t \geq 0}$ is a sequence of online gradient descent iterates on \mathcal{X} with step-sizes $(\gamma_t)_{t \geq 0}$, because for $0 \leq t \leq \tau - 1$, $u_t = 0$ and thus,

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t) = \Pi_{\mathcal{X}}(x_t + \gamma_t u_t),$$

and for $t \geq \tau$,

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t + \frac{\gamma}{\sqrt{\sum_{s=0}^t \|u_s\|_2^2}} u_t \right) = \Pi_{\mathcal{X}} (x_t + \gamma_t u_t).$$

Then the regret bound for OGD with time-dependent step-sizes from Corollary 3.3.16 gives

$$\begin{aligned} \sum_{t=0}^T \langle u_t, x - x_t \rangle &\leq \frac{\max_{0 \leq t \leq T} \|x - x_t\|_2^2}{2\gamma_T} + \sum_{t=0}^T \frac{\gamma_t \|u_t\|_2^2}{2} \\ &= \frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} \sqrt{\sum_{t=0}^T \|u_t\|_2^2} + \frac{\gamma}{2} \sum_{t=0}^T \frac{\|u_t\|_2^2}{\sqrt{\sum_{t=0}^T \|u_t\|_2^2}} \\ &\leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} + \gamma \right) \sqrt{\sum_{t=0}^T \|u_t\|_2^2}. \end{aligned}$$

using Lemma 7.2.1. Hence the result. \square

Remark 7.2.4.

Proposition 7.2.5 (Regret bound for AdaGrad-Diagonal). *Let $(u_t)_{t \geq 0}$ be a sequence in \mathbb{R}^d , $x_0 \in \mathcal{X}$, $\varepsilon, \gamma > 0$ and $(x_t)_{t \geq 0}$ the associated sequence of AdaGrad-Diagonal iterates on \mathcal{X} .*

(i) *Then for all $x \in \mathcal{X}$ and $T \geq 0$,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq \frac{\varepsilon}{2\gamma} \|x_0 - x\|_2^2 + \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_\infty^2}{2\gamma} + \gamma \right) \sum_{i=1}^d \sqrt{\sum_{t=0}^T u_{t,i}^2}.$$

(ii) *In particular, if $R \geq \max_{0 \leq t \leq T} \|x_t - x\|_\infty$, then $\gamma = R/\sqrt{2}$ yields,*

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq R \left(\frac{\varepsilon d}{\sqrt{2}} + \sum_{i=1}^d \sqrt{2 \sum_{t=0}^T u_{t,i}^2} \right).$$

(iii) *Moreover,*

$$\sum_{i=1}^d \sqrt{\sum_{t=0}^T u_{t,i}^2} = \inf_{\substack{v \in (\mathbb{R}_+^*)^d \\ \|v\|_1 \leq d}} \sqrt{d \sum_{t=0}^T \|u_t\|_{\text{diag}(v)^{-1}}^2} \leq \sqrt{d \sum_{t=0}^T \|u_t\|_2^2}.$$

Proof. Let $(A_t)_{t \geq 0}$ be the sequence of symmetric positive definite matrices defined as

$$A_t = \text{diag} \left(\varepsilon + \sqrt{\sum_{i=0}^t u_{t,i}^2} \right)_{1 \leq i \leq d},$$

and $(H_t)_{t \geq 0}$ the sequence of mirror maps defined as

$$H_t(x) = \frac{1}{2\gamma} \|x\|_{A_t}^2 = \frac{1}{2\gamma} \langle x, A_t x \rangle, \quad x \in \mathbb{R}^d, \quad t \geq 0.$$

Then, $(x_t)_{t \geq 0}$ corresponds to a sequence of online mirror descent iterates associated with mirror maps $(H_t)_{t \geq 0}$ and dual increments $(u_t)_{t \geq 0}$. Then Proposition 3.3.13 gives, with notation therein,

$$\sum_{t=0}^T \langle u_t, x - x_t \rangle \leq D_{H_0}(x, x_0) - D_{H_{T+1}}(x, x_{T+1}) + \sum_{t=0}^T \tilde{D}_{t+1/2}^\Delta + \sum_{t=0}^T \tilde{D}_t^*, \quad (7.1)$$

where for $t \geq 0$,

$$\tilde{D}_{t+1/2}^\Delta = D_{H_{t+1}-H_t}(x, x_{t+1}) = D_{H_{t+1}}(x, x_{t+1}) - D_{H_t}(x, x_{t+1}), \quad (7.2)$$

and since H_t is $1/\gamma$ -strongly convex for $\|\cdot\|_{A_t}$ by Corollary 1.6.9,

$$\tilde{D}_t^* = D_{H_t^*}(\nabla H_t(x_t) + u_t, \nabla H_t(x_t)) \leq \frac{\gamma}{2} \|u_t\|_{A_t^{-1}}^2.$$

Summing $\tilde{D}_{t+1/2}^\Delta$ and using the expression of the associated Bregman divergence from Example 1.5.7 gives

$$\begin{aligned} \sum_{t=0}^T \tilde{D}_{t+1/2}^\Delta &= \sum_{t=0}^T (D_{H_{t+1}}(x, x_{t+1}) - D_{H_t}(x, x_{t+1})) \\ &= D_{H_{T+1}}(x, x_{T+1}) - D_{H_0}(x, x_0) + \sum_{t=1}^T D_{H_t-H_{t-1}}(x, x_t) \end{aligned} \quad (7.3)$$

and the above last sum is bounded as

$$\begin{aligned}
\sum_{t=1}^T D_{H_t-H_{t-1}}(x, x_t) &= \frac{1}{2\gamma} \sum_{t=1}^T \|x - x_t\|_{A_t-A_{t-1}}^2 \\
&= \frac{1}{2\gamma} \sum_{t=1}^T \sum_{i=1}^d (x_i - x_{t,i})^2 (A_{t,ii} - A_{t-1,ii}) \\
&\leq \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \sum_{i=1}^d \sum_{t=1}^T (A_{t,ii} - A_{t-1,ii}) \\
&= \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \sum_{i=1}^d (A_{T,ii} - A_{0,ii}) \\
&= \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \sum_{i=1}^d \left(\varepsilon + \sqrt{\sum_{t=0}^T u_{t,i}^2} - \varepsilon - |u_{0,i}| \right) \\
&= \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \left(\sum_{i=1}^d \sqrt{\sum_{t=0}^T u_{t,i}^2} \right) \\
&\quad - \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \sum_{i=1}^d |u_{0,i}|.
\end{aligned} \tag{7.4}$$

Besides,

$$\begin{aligned}
D_H(x, x_0) &= \frac{1}{2\gamma} \|x - x_0\|_{A_0}^2 = \frac{1}{2\gamma} \sum_{i=1}^d (x_i - x_{0,i})^2 (\varepsilon + |u_{0,i}|) \\
&\leq \frac{\varepsilon}{2\gamma} \|x - x_0\|_2^2 + \frac{\max_{0 \leq t \leq T} \|x - x_t\|_\infty^2}{2\gamma} \sum_{i=1}^d |u_{0,i}|
\end{aligned} \tag{7.5}$$

Finally, summing \tilde{D}_t^* gives

$$\begin{aligned}
\sum_{t=0}^T \tilde{D}_t^* &\leq \frac{\gamma}{2} \sum_{t=0}^T \|u_t\|_{A_t^{-1}}^2 = \frac{\gamma}{2} \sum_{t=0}^T \sum_{i=1}^d \frac{u_{t,i}^2}{\varepsilon + \sqrt{\sum_{s=0}^t u_{s,i}^2}} \\
&\leq \frac{\gamma}{2} \sum_{t=0}^T \sum_{i=1}^d \frac{u_{t,i}^2}{\sqrt{\sum_{s=0}^t u_{s,i}^2}} \leq \gamma \sum_{i=1}^d \sqrt{\sum_{t=0}^T u_{t,i}^2},
\end{aligned} \tag{7.6}$$

where we used Lemma 7.2.1 for the last inequality.

Then combining (7.1), (7.2), (7.3), (7.4), (7.5) and (7.6) gives (i), and (ii) follows. Using Lemma 7.2.2 gives (iii). \square

7.3 Application to nonsmooth convex optimization

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex function that admits a minimizer \mathcal{X} , in other words, there exists $x_* \in \mathcal{X}$ such that

$$f(x_*) = \min_{x \in \mathcal{X}} f(x).$$

The following result demonstrates that AdaGrad-Norm is *adaptive* to the Lipschitz continuity of the objective function, meaning that it guarantees a convergence bound of order L/\sqrt{T} without prior knowledge of the Lipschitz coefficient L .

Proposition 7.3.1 (AdaGrad-Norm for nonsmooth convex optimization). *Assume that f is L -Lipschitz for $\|\cdot\|_2$. Let $R, \gamma > 0$, $x_0 \in \mathcal{X}$ and*

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t - \frac{\gamma}{\sum_{s=0}^t \|g_s\|_2^2} g_t \right), \quad t \geq 0,$$

where $g_t \in \partial f(x_t)$ for all $t \geq 0$.

(i) Then for all $T \geq 0$,

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} + \gamma \right) \frac{L}{\sqrt{T+1}}.$$

(ii) If $R \geq \max_{0 \leq t \leq T} \|x_t - x\|_2$, then $\gamma = R/\sqrt{2}$ yields

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq RL \sqrt{\frac{2}{T+1}}.$$

Proof. Combining the general regret bound for AdaGrad-Norm (Proposition 7.2.3) and Lemma 6.1.1 gives

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{1}{T+1} \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} + \gamma \right) \sqrt{\sum_{t=0}^T \|g_t\|_2^2}.$$

The Lipschitz continuity of f gives $\|g_t\|_2 \leq L$ for all $t \geq 0$. The result follows. \square

The following result demonstrates that, in the context of stochastic convex optimization, AdaGrad-Norm is *adaptive* to the second-order moment of the gradient estimators: if the latter are bounded by σ^2 , it achieves a convergence bound of order σ/\sqrt{T} without prior knowledge of σ .

Proposition 7.3.2 (AdaGrad-Norm for stochastic nonsmooth convex optimization). *Let $R, \sigma > 0$, $x_0 \in \mathcal{X}$ and*

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t - \frac{R}{\sqrt{2 \sum_{s=0}^t \|g_s\|_2^2}} g_t \right), \quad t \geq 0,$$

where for each $t \geq 0$, almost-surely,

$$\mathbb{E}[g_t | x_t] \in \partial f(x_t), \quad \mathbb{E}[\|g_t\|_2^2 | x_t] \leq \sigma^2, \quad \text{and} \quad \|x_t - x_*\|_2 \leq R.$$

Then, for all $T \geq 0$,

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq R\sigma \sqrt{\frac{2}{T+1}}.$$

Proof. Combining the general regret bound for AdaGrad-Norm (Proposition 7.2.3) and Lemma 6.4.2 gives

$$\mathbb{E} \left[\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right] \leq \frac{R}{T+1} \mathbb{E} \left[\sqrt{2 \sum_{t=0}^T \|g_t\|_2^2} \right].$$

Then using Jensen's inequality with the concavity of the square root,

$$\begin{aligned} \mathbb{E} \left[\sqrt{2 \sum_{t=0}^T \|g_t\|_2^2} \right] &\leq \sqrt{2 \cdot \mathbb{E} \left[\sum_{t=0}^T \|g_t\|_2^2 \right]} \\ &\leq \frac{R}{T+1} \mathbb{E} \left[\sqrt{2 \sum_{t=0}^T \mathbb{E}[\|g_t\|_2^2 | x_t]} \right] \\ &= R\sigma \sqrt{\frac{2}{T+1}}, \end{aligned}$$

hence the result. \square

7.4 Application to smooth convex optimization

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex function that is L -smooth for $\|\cdot\|_2$ and that admits a global minimizer x_* :

$$f(x_*) = \min_{x \in \mathbb{R}^d} f(x).$$

Lemma 7.4.1. *For all $x \in \mathbb{R}^d$,*

$$\|\nabla f(x)\|_2^2 \leq 2L(f(x) - f(x_*)).$$

Proof. Let $x' \in \mathbb{R}^d$. The definition of smoothness gives

$$f(x') - f(x) - \langle \nabla f(x), x' - x \rangle \leq \frac{L}{2} \|x' - x\|_2^2.$$

For $x' = x - \frac{1}{L} \nabla f(x)$, the above simplifies into

$$f(x') - f(x) + \frac{1}{2L} \|\nabla f(x)\|_2^2.$$

The result follows because $f(x_*) \leq f(x')$. \square

The following statement proves that AdaGrad-Norm is *adaptive* to the smoothness of the objective function: it achieves a L/T convergence bound without prior knowledge of the smoothness coefficient L .

Proposition 7.4.2 (AdaGrad-Norm for smooth convex optimization). *Assume $x_* \in \mathcal{X}$. Let $\gamma > 0$, $x_0 \in \mathcal{X}$ and*

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t - \frac{\gamma}{\sum_{s=0}^t \|\nabla f(x_s)\|_2^2} \nabla f(x_t) \right), \quad t \geq 0.$$

(i) *Let $T \geq 0$. Then,*

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right)^2 \frac{2L}{T+1}.$$

(ii) *Let $R \geq \max_{0 \leq t \leq T} \|x_t - x_*\|_2$, then $\gamma = R/\sqrt{2}$ yields,*

$$\min_{0 \leq t \leq T} f(x_t) - f(x_*) \leq \frac{4R^2L}{T+1}.$$

Proof. Using general regret bound for AdaGrad-Norm (Proposition 7.2.3) and the convexity of f ,

$$\begin{aligned} \sum_{t=0}^T (f(x_t) - f(x_*)) &\leq \sum_{t=0}^T \langle \nabla f(x_t), x_t - x_* \rangle \\ &\leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right) \sqrt{\sum_{t=0}^T \|\nabla f(x_t)\|_2^2} \\ &\leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right) \sqrt{2L \sum_{t=0}^T (f(x_t) - f(x_*))}, \end{aligned}$$

where we used Lemma 7.4.1 for the last inequality. Dividing by $\sqrt{\sum_{t=0}^T (f(x_t) - f(x_*))}$ and taking the square gives

$$\sum_{t=0}^T (f(x_t) - f(x_*)) \leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x\|_2^2}{2\gamma} + \gamma \right)^2 2L.$$

The above left-hand side is bounded from below as

$$\sum_{t=0}^T (f(x_t) - f(x_*)) \geq \sum_{t=0}^T \left(\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right) \geq T \left(\min_{0 \leq t \leq T} f(x_t) - f(x_*) \right),$$

hence the result. \square

Chapter 8

Monotone operators and fixed point iterations

This chapter gives a quick preview of monotone operators, fixed point iterations, and their relation to regret minimization. Monotone operators are a generalization of the gradient of a convex function, and allow to deal with various problems such as convex-concave saddle-points, convex games, finite two-player zero-sum games, etc.

Monotone operators can be defined with set-valued mappings, to generalize subdifferentials, but we restrict to single-valued mappings for simplicity.

Let $\mathcal{X} \subset \mathbb{R}^d$ be nonempty closed and convex. I denotes the identity map on a set which will be clear from the context.

8.1 Monotone operators

Recall the minimization of a differentiable convex function $f : \mathbb{R}^d \rightarrow \mathbb{R}$. A global solution corresponds to a *zero* of the gradient: the corresponding notion below is the zero of the monotone operator. Regarding the constrained problem on \mathcal{X} , according to Proposition 1.2.10, the minimizer x_* of f on \mathcal{X} is characterized by the following variational inequality:

$$\forall x \in \mathcal{X}, \quad \langle \nabla f(x_*), x - x_* \rangle \geq 0,$$

which below is extended to the notion of *strong solution* on \mathcal{X} of a monotone operator.

Definition 8.1.1. Let $G : \mathcal{X} \rightarrow \mathbb{R}^d$ and $x_* \in \mathcal{X}$.

(i) G is a *monotone operator* if:

$$\forall x, x' \in \mathcal{X}, \quad \langle G(x') - G(x), x' - x \rangle \geq 0.$$

(ii) x_* is a zero of G if $G(x_*) = 0$.

(iii) x_* is a *strong solution* of G on \mathcal{X} if

$$\forall x \in \mathcal{X}, \quad \langle G(x_*), x_* - x \rangle \leq 0.$$

(iv) x_* is a *weak solution* of G on \mathcal{X} if

$$\forall x \in \mathcal{X}, \quad \langle G(x), x_* - x \rangle \leq 0.$$

Remark 8.1.2. An advantage of the concept of weak solution is that such a solution always exist as soon as \mathcal{X} is compact.

Remark 8.1.3. If x_* is either a weak or a strong solution and belongs to the interior of \mathcal{X} , then it is a zero of the operator.

The following proposition proves that in the case of a continuous monotone operator, both above solution concepts are equivalent.

Proposition 8.1.4. *Let $G : \mathcal{X} \rightarrow \mathbb{R}^d$ a monotone operator and $x_* \in \mathcal{X}$.*

(i) *If x_* is a strong solution of G on \mathcal{X} , then it is a weak solution of G on \mathcal{X} .*

(ii) *If x_* is a weak solution of G on \mathcal{X} and G is continuous, then it is a strong solution of G on \mathcal{X} .*

Proof. (i) If x_* is a strong solution, for all $x \in \mathcal{X}$,

$$\langle G(x), x_* - x \rangle \leq \langle G(x_*), x_* - x \rangle \leq 0,$$

where the first inequality holds by monotonicity of G .

(ii) Now assume that x_* is a weak solution and that G is continuous. Let us prove that x_* is a strong solution. Let $x \in \mathcal{X}$ and for all $\lambda \in (0, 1)$ consider $x_\lambda = (1 - \lambda)x_* + \lambda x$, which belongs to \mathcal{X} by convexity of the latter. Then because x_* is a weak solution,

$$\langle G(x_\lambda), x_* - x_\lambda \rangle \leq 0,$$

where $x_* - x_\lambda$ rewrites as $\lambda(x_* - x)$. Then, dividing by λ gives

$$\langle G(x_\lambda), x_* - x \rangle \leq 0.$$

Taking the limit as $\lambda \rightarrow 0^+$ gives, by continuity of G ,

$$\langle G(x_*), x_* - x \rangle \leq 0.$$

x_* is thus a strong solution. □

Proposition 8.1.5. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a differentiable convex function. Then, $\nabla f : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a monotone operator.*

Proof. Let $x, x' \in \mathbb{R}^d$. By convexity,

$$\begin{aligned} f(x') &\geq f(x) + \langle \nabla f(x), x' - x \rangle \\ f(x) &\geq f(x') + \langle \nabla f(x'), x - x' \rangle. \end{aligned}$$

Summing the above two inequalities gives

$$\langle \nabla f(x') - \nabla f(x), x' - x \rangle \geq 0,$$

hence the result. \square

Definition 8.1.6. Let $m, n \geq 1$ be integers, $g : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$, $\mathcal{A} \subset \mathbb{R}^m$ and $\mathcal{B} \subset \mathbb{R}^n$ nonempty sets. A couple $(a_*, b_*) \in \mathcal{A} \times \mathcal{B}$ is a *saddle-point* of g on $\mathcal{A} \times \mathcal{B}$ if

$$a_* \in \underset{a \in \mathcal{A}}{\text{Arg min}} g(a, b_*) \quad \text{and} \quad b_* \in \underset{b \in \mathcal{B}}{\text{Arg max}} g(a_*, b).$$

If moreover $\mathcal{A} = \mathbb{R}^m$ and $\mathcal{B} = \mathbb{R}^n$, the saddle-point is said to be *global*.

Proposition 8.1.7. Let $m, n \geq 1$ be integer, $g : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- for all $b \in \mathbb{R}^n$, $g(\cdot, b)$ is convex and differentiable (denote $\nabla_{ag}(\cdot, b)$ its gradient);
- for all $a \in \mathbb{R}^m$, $g(a, \cdot)$ is concave and differentiable (denote $\nabla_{bg}(a, \cdot)$ its gradient).

Let $G : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^m \times \mathbb{R}^n$ be defined as

$$G(a, b) = (\nabla_{ag}(a, b), -\nabla_{bg}(a, b)), \quad a \in \mathbb{R}^m, b \in \mathbb{R}^n,$$

and $\mathcal{A} \subset \mathbb{R}^m$ and $\mathcal{B} \subset \mathbb{R}^n$ be nonempty closed convex sets.

(i) G is a monotone operator.

(ii) $(a, b) \in \mathcal{A} \times \mathcal{B}$ is a saddle-point of g on $\mathcal{A} \times \mathcal{B}$ if, and only if it is a strong solution of G on $\mathcal{A} \times \mathcal{B}$.

(iii) $(a, b) \in \mathbb{R}^m \times \mathbb{R}^n$ is a global saddle-point of g if, and only if, it is a zero of G .

Example 8.1.8 (KKT operator). Let $p \geq 1$ be an integer, $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a differentiable convex function, $A \in \mathbb{R}^{p \times d}$ and $b \in \mathbb{R}^p$. Consider the constrained optimization problem

$$\begin{aligned} &\text{minimize} && f(x) \\ &\text{subject to} && Ax = b. \end{aligned}$$

Solving the above is equivalent to finding a saddle-point of the Lagrangian $L(x, w) = f(x) + w^\top(Ax - b)$, which in turn is equivalent to finding a zero of the associated KKT operator:

$$F(x, w) = \begin{pmatrix} \nabla f(x) + A^\top w \\ b - Ax \end{pmatrix}.$$

Proposition 8.1.9. *The Euclidean projection on a nonempty closed convex subset of \mathbb{R}^d is a monotone operator.*

8.2 Bounded monotone operators

We consider bounded monotone operators, which is the natural extension of Lipschitz convex optimization. The simplest algorithm for this problem is the obvious extension of projected gradient descent, which is called the (projected) *forward step* algorithm:

$$x_{t+1} = \Pi_{\mathcal{X}}(x_t - \gamma_t G(x_t)), \quad t \geq 0.$$

For a class of UMD-based iterates that contains the above, Proposition 8.2.3 guarantees that an approximate weak solution can be obtained at speed $1/\sqrt{T}$. Then, in the special case of AdaGrad-Norm step-sizes, Proposition 7.2.5 provides a guarantee that is *adaptive* to the bound on the operator.

Let $G : \mathcal{X} \rightarrow \mathbb{R}^d$ be a monotone operator.

Example 8.2.1 (Finite two-player zero-sum games). Let $m, n \geq 1$ be integers, and $A \in \mathbb{R}^{m \times n}$. The solutions of the corresponding two-player zero-sum games are the solutions of the following constrained saddle-point problem:

$$\max_{a \in \Delta_m} \min_{b \in \Delta_n} \langle a, Ab \rangle.$$

The corresponding monotone operator $G(a, b) = (-Ab, A^\top a)$ is bounded on $\Delta_m \times \Delta_n$.

The following gives a connection between the search for an approximate weak solution and regret minimization.

Lemma 8.2.2. *Let $(x_t)_{t \geq 0}$ be a sequence in \mathcal{X} , $(\gamma_t)_{t \geq 0}$ a positive sequence, $T \geq 0$, and*

$$\bar{x}_T^{(\gamma)} = \frac{\sum_{t=0}^T \gamma_t x_t}{\sum_{t=0}^T \gamma_t}.$$

Then for all $x \in \mathcal{X}$,

$$\langle G(x), \bar{x}_T^{(\gamma)} - x \rangle \leq \left(\sum_{t=0}^T \gamma_t \right)^{-1} \sum_{t=0}^T \langle \gamma_t G(x_t), x_t - x \rangle.$$

Proof. For all $t \geq 0$, the monotonicity of G gives

$$\gamma_t \langle G(x), x_t - x \rangle \leq \gamma_t \langle G(x_t), x_t - x \rangle.$$

Summing and dividing by $\sum_{t=0}^T \gamma_t$ gives the result. \square

Proposition 8.2.3 (Strict UMD iterates with time-dependent step-sizes and non-uniform averaging for variational inequalities with bounded monotone operators). *Let $\|\cdot\|$ be a norm on \mathbb{R}^d , $K, R > 0$, and h a regularizer on \mathcal{X} . We assume that*

- $(x_t, y_t)_{t \geq 0}$ is a sequence of strict UMD iterates associated with regularizer h and dual increments $(-\gamma_t G(x_t))_{t \geq 0}$,
- h is K -strongly convex for $\|\cdot\|$,
- for all $x \in \mathcal{X}$, $\|G(x)\|_* \leq L$.

(i) Let $T \geq 0$, $x \in \text{dom } h$ and $\bar{x}_T^{(\gamma)} = \left(\sum_{t=0}^T \gamma_t\right)^{-1} \sum_{t=0}^T \gamma_t x_t$. Then,

$$\forall x \in \text{dom } h, \quad \langle G(x), \bar{x}_T^{(\gamma)} - x \rangle \leq \frac{D_h(x, x_0; y_0) + \frac{L^2}{2K} \sum_{t=0}^T \gamma_t^2}{\sum_{t=0}^T \gamma_t}.$$

(ii) Let $\gamma > 0$. If $\gamma_t = \gamma \sqrt{2K} / (L\sqrt{t+1})$ for all $t \geq 0$, then

$$\langle G(x), \bar{x}_T^{(\gamma)} - x \rangle \leq \frac{L}{\sqrt{2KT}} \left(\frac{D_h(x, x_0; y_0)}{\gamma} + \gamma(1 + \log(T+1)) \right).$$

(iii) If $D_h(x, x_0; y_0) \leq R^2$, then $\gamma = R$ yields,

$$\langle G(x), \bar{x}_T^{(\gamma)} - x \rangle \leq \frac{RL}{\sqrt{2KT}} (2 + \log(T+1)).$$

Proof. Use above Lemma 8.2.2 and perform a similar analysis as for Lipschitz convex optimization (Proposition 6.1.2). \square

From the above general statement, we can of course derive corollaries for OMD and DA. Analogously to Proposition 6.1.6, DA with time-dependent parameters can be used instead to shave off the $\log T$ factor.

Proposition 8.2.4 (AdaGrad-Norm for bounded monotone operators). *Assume that G is bounded by L with respect to $\|\cdot\|_2$. Let $x_0 \in \mathcal{X}$, $\gamma > 0$,*

$$x_{t+1} = \Pi_{\mathcal{X}} \left(x_t - \frac{\gamma}{\sqrt{\sum_{s=0}^t \|G(x_s)\|_2^2}} G(x_t) \right), \quad t \geq 0.$$

(i) Let $T \geq 0$ and $x \in \mathcal{X}$. Then,

$$\langle G(x), \bar{x}_T - x \rangle \leq \left(\frac{\max_{0 \leq t \leq T} \|x - x_t\|_2^2}{2\gamma} + \gamma \right) \sqrt{\frac{L}{T+1}},$$

where $\bar{x}_T = \frac{1}{T+1} \sum_{t=0}^T x_t$.

(ii) Let $R \geq \max_{0 \leq t \leq T} \|x_t - x\|_2$, then $\gamma = R/\sqrt{2}$ yields

$$\langle G(x), \bar{x}_T - x \rangle \leq RL \sqrt{\frac{2}{T+1}}.$$

Proof. Combine above Lemma 8.2.2 with the regret bound for AdaGrad-Norm (Proposition 7.2.3). \square

8.3 Lipschitz continuous monotone operators

We now consider Lipschitz continuous monotone operators, for which approximate weak solutions can be obtained at rate $1/T$ instead of $1/\sqrt{T}$. An important special case discussed below is two-player zero-sum games. The simplest method with such a guarantee is the *extragradient* algorithm which writes

$$\begin{aligned} w_t &= \Pi_{\mathcal{X}}(x_t - \gamma G(x_t)), \\ x_{t+1} &= \Pi_{\mathcal{X}}(x_t - \gamma G(w_t)), \quad t \geq 0, \end{aligned}$$

where $\gamma > 0$ is a step-size. We present and analyze below a very general a class of UMD-based iterates that extends extragradient.

Example 8.3.1 (Finite two-player zero-sum games). Let $m, n \geq 1$ be integers, and $A \in \mathbb{R}^{m \times n}$. Corresponding monotone operator $G(a, b) = (-Ab, A^\top a)$ is Lipschitz continuous on $\Delta_m \times \Delta_n$.

Let $G : \mathcal{X} \rightarrow \mathbb{R}^d$ be a monotone operator.

Definition 8.3.2. Let h a regularizer on \mathcal{X} and $\gamma > 0$. A sequence $((x_t, w_t, y_t, z_t))_{t \geq 0}$ in $(\mathbb{R}^d)^4$ is a sequence of *UMP iterates* associated with regularizer h , operator G and step-size γ if $((x_t, y_t))_{t \geq 0}$ is a sequence of strict UMD iterates associated with regularizer h and dual iterates $(-\gamma G(w_t))_{t \geq 0}$ and for $t \geq 0$,

- (i) $z_t \in \partial h(x_t)$,
- (ii) $\forall x \in \mathcal{X}, \langle z_t - y_t, x - x_t \rangle \geq 0$,
- (iii) $w_t = \nabla h^*(z_t - \gamma G(x_t))$,

Proposition 8.3.3 (UMP iterates for variational inequalities with Lipschitz continuous monotone operator). *Let $K, L > 0$, $\|\cdot\|$ be a norm on \mathbb{R}^d , h a regularizer on \mathcal{X} , $((x_t, y_t, w_t, z_t))_{t \geq 0}$ a sequence of UMP iterates associated with regularizer h , operator G and step-size K/L . We assume that*

- h is K -strongly convex for $\|\cdot\|$,

- $G : \mathcal{X} \rightarrow \mathbb{R}^d$ is L -Lipschitz continuous for the following norms:

$$\|G(x') - G(x)\|_* \leq L \|x' - x\|, \quad x, x' \in \mathcal{X}.$$

Then for all $T \geq 0$,

$$\forall x \in \text{dom } h, \quad \langle G(x), \bar{w}_T - x \rangle \leq \frac{L \cdot D_h(x, x_0; y_0)}{K(T+1)},$$

where $\bar{w}_T = \frac{1}{T+1} \sum_{t=0}^T w_t$.

Proof. Let $x \in \text{dom } h$ and $T \geq 0$. Lemma 8.2.2 gives:

$$\begin{aligned} \frac{K(T+1)}{L} \langle G(x) | \bar{w}_T - x \rangle &\leq \sum_{t=0}^T \frac{K}{L} \langle G(w_t) | w_t - x \rangle \\ &= \sum_{t=0}^T \left(\frac{K}{L} \langle G(w_t) | x_{t+1} - x \rangle + \frac{K}{L} \langle G(w_t) | w_t - x_{t+1} \rangle \right) \\ &\leq D_h(x, x_0; y_0) + \sum_{t=0}^T \left(-D_h(x_{t+1}, x_t; y_t) + \frac{K}{L} \langle G(w_t) | w_t - x_{t+1} \rangle \right), \end{aligned}$$

where the second inequality comes from applying Lemma 2.4.1 (because $(x_t, y_t)_{t \geq 1}$ is a sequence of strict UMD iterates). We bound the above last two terms as follows. Let $t \geq 0$ and denote δ_t the content of the above last sum and let us bound it as follows.

$$\begin{aligned} \delta_t &= -D_h(x_{t+1}, x_t; y_t) - \frac{K}{L} \langle G(w_t) | x_{t+1} - w_t \rangle \\ &= -h(x_{t+1}) + h(x_t) + \langle y_t | x_{t+1} - x_t \rangle + \frac{K}{L} \langle (G(x_t) - G(w_t)) | x_{t+1} - w_t \rangle \\ &\quad - \frac{K}{L} \langle G(x_t) | x_{t+1} - w_t \rangle. \end{aligned} \tag{8.1}$$

Condition (ii) from the Definition 8.3.2:

$$\langle y_t | x_{t+1} - x_t \rangle \leq \langle z_t | x_{t+1} - x_t \rangle. \tag{8.2}$$

Besides, using basic inequality $\langle y | x \rangle \leq \frac{1}{2} \|y\|_*^2 + \frac{1}{2} \|x\|^2$, we can write:

$$\begin{aligned} \frac{K}{L} \langle (G(x_t) - G(w_t)) | x_{t+1} - w_t \rangle &\leq \frac{(K/L)^2}{2K} \|G(x_t) - G(w_t)\|_*^2 + \frac{K}{2} \|x_{t+1} - w_t\|^2 \\ &\leq \frac{(K/L)^2 L^2}{2K} \|x_t - w_t\|^2 + \frac{K}{2} \|x_{t+1} - w_t\|^2, \end{aligned} \tag{8.3}$$

where we used the Lipschitz continuity of operator G . Injecting (8.2) and (8.3) into (8.1) and simplifying gives

$$\begin{aligned} \delta_t \leq & -h(x_{t+1}) + h(x_t) + \langle z_t | w_t - x_t \rangle + \left\langle z_t - \frac{K}{L}G(x_t) \middle| x_{t+1} - w_t \right\rangle \\ & + \frac{K}{2} \|w_t - x_t\|_*^2 + \frac{K}{2} \|x_{t+1} - w_t\|^2. \end{aligned} \quad (8.4)$$

First, we have $z_t \in \partial h(x_t)$ by condition (i) and $D_h(w_t, x_t; z_t)$ is well-defined. Besides, thanks to Proposition 1.4.6, condition (iii) is equivalent to $z_t - \frac{K}{L}G(x_t) \in \partial h(w_t)$, and $D_h(x_{t+1}, w_t; z_t - \frac{K}{L}G(x_t))$ is thus well-defined. We can make those two generalized Bregman divergences appear in the above right-hand side, which is consequently equal to:

$$\begin{aligned} \delta_t \leq & \frac{K}{2} \|w_t - x_t\|^2 - D_h(w_t, x_t; z_t) + \frac{K}{2} \|x_{t+1} - w_t\|^2 \\ & - D_h\left(x_{t+1}, w_t; z_t - \frac{K}{L}G(x_t)\right). \end{aligned}$$

Using the K -strong convexity of h (Proposition 1.6.4), the above simplifies to $\delta_t \leq 0$. The result follows. \square

The above general statement contains as special cases the following well-known algorithms.

Corollary 8.3.4 (Mirror-Prox). *Let $K, L > 0$, $\|\cdot\|$ be a norm on \mathbb{R}^d , H a mirror map compatible with \mathcal{X} , $x_0 \in \mathcal{X} \cap \text{int dom } H$ and for $t \geq 0$,*

$$\begin{aligned} w_t &= \arg \max_{x \in \mathcal{X}} \left\{ \left\langle \nabla H(x_t) - \frac{K}{L}G(x_t), x \right\rangle - H(x) \right\} \\ x_{t+1} &= \arg \max_{x \in \mathcal{X}} \left\{ \left\langle \nabla H(x_t) - \frac{K}{L}G(w_t), x \right\rangle - H(x) \right\}. \end{aligned}$$

Assume that

- H is K -strongly convex for $\|\cdot\|$,
- $G : \mathcal{X} \rightarrow \mathbb{R}^d$ is L -Lipschitz continuous for the following norms:

$$\|G(x') - G(x)\|_* \leq L \|x' - x\|, \quad x, x' \in \mathcal{X}.$$

Then for all $T \geq 0$,

$$\forall x \in \mathcal{X} \cap \text{dom } H, \quad \langle G(x), \bar{w}_T - x \rangle \leq \frac{L \cdot D_H(x, x_0)}{K(T+1)},$$

where $\bar{w}_T = \frac{1}{T+1} \sum_{t=0}^T w_t$.

Corollary 8.3.5 (Extragradient). *Let $L > 0$, $x_0 \in \mathcal{X}$ and for $t \geq 0$,*

$$\begin{aligned} w_t &= \Pi_{\mathcal{X}} \left(x_t - \frac{1}{L} G(x_t) \right) \\ x_{t+1} &= \Pi_{\mathcal{X}} \left(x_t - \frac{1}{L} G(w_t) \right). \end{aligned}$$

Assume that G is L -Lipschitz continuous for $\|\cdot\|_2$. Then, for all $T \geq 0$,

$$\forall x \in \mathcal{X}, \quad \langle G(x), \bar{w}_T - x \rangle \leq \frac{L \|x - x_0\|_2^2}{2(T+1)},$$

where $\bar{w}_T = \frac{1}{T+1} \sum_{t=0}^T w_t$.

Proof. Apply Corollary 8.3.4 with the Euclidean mirror map. \square

Corollary 8.3.6 (Dual extrapolation). *Under the assumptions of Corollary 8.3.4, let $h = H + I_{\mathcal{X}}$, and for $t \geq 0$,*

$$\begin{aligned} x_t &= \nabla h^* \left(y_0 - \frac{K}{L} \sum_{s=0}^{t-1} G(w_s) \right), \\ w_t &= \arg \max_{x \in \mathcal{X}} \left\{ \left\langle \nabla H(x_t) - \frac{K}{L} G(x_t), x \right\rangle - H(x) \right\}. \end{aligned}$$

Then for all $T \geq 0$,

$$\forall x \in \text{dom } h, \quad \langle G(x), \bar{w}_T - x \rangle \leq \frac{L \cdot D_h(x, x_0; y_0)}{K(T+1)},$$

where $\bar{w}_T = \frac{1}{T+1} \sum_{t=0}^T w_t$.

8.4 Co-coercive operators and fixed point iterations

We now consider co-coercivity, which is a stronger notion than monotonicity. We establish the equivalence between being a zero of a co-coercive operator and being a fixed point of a corresponding *nonexpansive* map (meaning 1-Lipschitz continuous for the Euclidean norm). This subtopic is particularly important as it covers the construction and analysis of many optimization algorithms, such as the proximal gradient descent, ADMM, Douglas-Rachford splitting, Chambolle-Pock, and many more.

We present in Lemma 8.4.8 a connection between the search for an approximate solution of a co-coercive operator and regret minimization. We use this connection to recover in Proposition 8.4.9 the classical guarantee

for the Krasnoselskii-Mann method, which is the workhorse fixed point iteration for nonexpansive maps. We further utilize this connection to define an AdaGrad-Norm-based method for fixed points which comes in Proposition 8.4.10 with an *adaptive* guarantee.

Definition 8.4.1. Let $L > 0$. A map $G : \mathcal{X} \rightarrow \mathbb{R}^d$ is a *L-co-coercive* operator if for all $x, x' \in \mathcal{X}$,

$$\langle G(x') - G(x), x' - x \rangle \geq \frac{1}{L} \|G(x') - G(x)\|_2^2.$$

Definition 8.4.2. A map $F : \mathcal{X} \rightarrow \mathcal{X}$ is *nonexpansive* if it is 1-Lipschitz continuous for $\|\cdot\|_2$, in other words

$$\forall x, x' \in \mathcal{X}, \quad \|F(x') - F(x)\|_2 \leq \|x' - x\|_2.$$

Definition 8.4.3. Let $F : \mathcal{X} \rightarrow \mathcal{X}$. x_* is a fixed point of F if $F(x_*) = x_*$.

Proposition 8.4.4. Let $F : \mathcal{X} \rightarrow \mathcal{X}$, $x_* \in \mathcal{X}$ and $L > 0$. The following statements are equivalent.

- (i) x_* is a fixed point of F ,
- (ii) x_* is a fixed point of $\frac{1}{L}F + (1 - \frac{1}{L})I$,
- (iii) x_* is a zero of $I - F$.

Proof. Immediate. □

Proposition 8.4.5. Let $L > 0$. A map $G : \mathcal{X} \rightarrow \mathbb{R}^d$ is a *L-co-coercive* operator if, and only if, $I - \frac{2}{L}G$ is nonexpansive.

Proof. For $G : \mathcal{X} \rightarrow \mathbb{R}^d$, being *L-co-coercive* can be written, for all $x, x' \in \mathcal{X}$,

$$\frac{1}{L} \|G(x') - G(x)\|_2^2 - \langle G(x') - G(x), x' - x \rangle \leq 0.$$

Multiplying by $4/L$ and adding $\|x - x'\|_2^2$, the above equivalently rewrites: for all $x, x' \in \mathcal{X}$,

$$\begin{aligned} \left\| \frac{2}{L}G(x') - \frac{2}{L}G(x) \right\|_2^2 - 2 \left\langle \frac{2}{L}G(x') - \frac{2}{L}G(x), x' - x \right\rangle + \|x' - x\|_2^2 \\ \leq \|x' - x\|_2^2, \end{aligned}$$

which simplifies into

$$\left\| \left(\frac{2}{L}G - I \right) (x') - \left(\frac{2}{L}G - I \right) (x) \right\|_2^2 \leq \|x' - x\|_2^2,$$

in other words, $I - \frac{2}{L}G$ is nonexpansive. □

Corollary 8.4.6. *Let $L > 0$ and $F : \mathcal{X} \rightarrow \mathcal{X}$. Then, $\frac{1}{L}F + (1 - \frac{1}{L})I$ is nonexpansive if, and only if $\frac{1}{2}(I - F)$ is L -co-coercive.*

Proof. Follows from Proposition 8.4.5. \square

Proposition 8.4.7. *Let $L > 0$ and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ a differentiable convex function. Then, f is L -smooth if, and only if, ∇f is a L -co-coercive operator.*

Proof. This characterization is part of Proposition 1.6.4. \square

The easiest fixed points problems are those with a contraction map $F : \mathcal{X} \rightarrow \mathcal{X}$ (meaning L -Lipschitz continuous for $\|\cdot\|_2$ with $0 \leq L < 1$). Banach's fixed point theorem states that, from any initial point $x_0 \in \mathcal{X}$, iteration $x_{t+1} = F(x_t)$ ensures *geometric convergence* to the fixed point, which is necessarily unique.

When the map is only nonexpansive, the above iteration does not ensure convergence anymore. This motivates the Krasnoselskii-Mann iteration (see Proposition 8.4.9 below) which is a damped version: the next iterate x_{t+1} is obtained by the convex combination of the image $F(x_t)$ with the current iterate x_t . Convergence to a fixed point is then ensured for e.g. constant coefficients (for the convex combination), but there no more guarantees on the distance to the fixed point. Instead, a vanishing bound on $\|F(x_t) - x_t\|$ is obtained, which is a weaker guarantee but somehow still measures how far x_t is from being a fixed point.

Lemma 8.4.8. *Let $L > 0$, $F : \mathcal{X} \rightarrow \mathcal{X}$ such that $\frac{1}{L}F + (1 - \frac{1}{L})I$ is nonexpansive, x_* a fixed point of F , $(x_t)_{t \geq 0}$ be a sequence in \mathcal{X} , $(\gamma_t)_{t \geq 0}$ a positive sequence, and $T \geq 0$. Then,*

$$\sum_{t=0}^T \gamma_t \|F(x_t) - x_t\|_2^2 \leq 2L \sum_{t=0}^T \langle \gamma_t (F(x_t) - x_t), x_* - x_t \rangle.$$

Proof. For $t \geq 0$, because $\frac{1}{2}(I - F)$ is L -co-coercive by Corollary 8.4.6,

$$\begin{aligned} \gamma_t \langle F(x_t) - x_t, x_* - x_t \rangle &= 2\gamma_t \left\langle \frac{F(x_t) - x_t}{2} - \frac{F(x_*) - x_*}{2}, x_* - x_t \right\rangle \\ &\geq \frac{2\gamma_t}{L} \left\| \frac{F(x_t) - x_t}{2} - \frac{F(x_*) - x_*}{2} \right\|_2^2 \\ &= \frac{\gamma_t}{2L} \|F(x_t) - x_t\|_2^2. \end{aligned}$$

The result follows by summing the weighted average. \square

Proposition 8.4.9 (Krasnoselskii-Mann iterations). *Let $F : \mathcal{X} \rightarrow \mathcal{X}$ be a nonexpansive map, x_* a fixed point of F , $(\gamma_t)_{t \geq 0}$ a sequence in $(0, 1)$, $x_0 \in \mathcal{X}$ and for $t \geq 0$,*

$$x_{t+1} = (1 - \gamma_t)x_t + \gamma_t F(x_t).$$

(i) $(x_t)_{t \geq 0}$ is a sequence of OGD iterates on \mathbb{R}^d associated with dual increments $(\gamma_t(F(x_t) - x_t))_{t \geq 0}$.

(ii) Let $T \geq 0$. Then,

$$\|F(x_T) - x_T\|_2 \leq \frac{\|x_0 - x_*\|_2}{\sqrt{\sum_{t=0}^T \gamma_t(1 - \gamma_t)}}.$$

(iii) In particular, if $\gamma_t = 1/2$ for all $t \geq 0$,

$$\|F(x_T) - x_T\|_2 \leq \frac{2\|x_0 - x_*\|_2}{\sqrt{T+1}}.$$

Proof. For $t \geq 0$, the definition of the iterates rewrites

$$x_{t+1} = x_t + \gamma_t(F(x_t) - x_t),$$

which gives (i).

Combining the regret bound for OGD (Corollary 3.3.16) with Lemma 8.4.8 gives

$$\sum_{t=0}^T \gamma_t \|F(x_t) - x_t\|_2^2 \leq \|x_0 - x_*\|_2^2 - \|x_{T+1} - x_*\|_2^2 + \sum_{t=0}^T \gamma_t^2 \|F(x_t) - x_t\|_2^2.$$

Getting rid of term $\|x_{T+1} - x_*\|_2^2$ and rearranging gives

$$\sum_{t=0}^T \gamma_t(1 - \gamma_t) \|F(x_t) - x_t\|_2^2 \leq \|x_0 - x_*\|_2^2.$$

The result follows as soon as $(\|F(x_t) - x_t\|_2^2)_{t \geq 0}$ is nonincreasing, which we now prove. Let $t \geq 0$. Note that by definition of the iterates, $F(x_t) - x_{t+1} = (1 - \gamma_t)(F(x_t) - x_t)$. Then,

$$\begin{aligned} \|F(x_{t+1}) - x_{t+1}\|_2 &= \|F(x_{t+1}) - F(x_t) + F(x_t) - x_{t+1}\|_2 \\ &\leq \|F(x_{t+1}) - F(x_t)\|_2 + \|F(x_t) - x_{t+1}\|_2 \\ &\leq \|x_{t+1} - x_t\|_2 + (1 - \gamma_t) \|F(x_t) - x_t\|_2 \\ &= \gamma_t \|F(x_t) - x_t\|_2 + (1 - \gamma_t) \|F(x_t) - x_t\|_2, \end{aligned}$$

where we used the nonexpansiveness of F for the second inequality. Hence (ii) and (iii) follows. \square

We now transpose the AdaGrad-Norm algorithm to the problem of finding a fixed point of a map F , which may not be nonexpansive, but such that $\frac{1}{L}F + (1 - \frac{1}{L})I$ is nonexpansive, for some $L > 0$. The adaptive character of AdaGrad allows to be adaptive to such a coefficient L , without prior knowledge.

Proposition 8.4.10 (AdaGrad-Norm for fixed points). *Let $L, R > 0$, $F : \mathcal{X} \rightarrow \mathcal{X}$ such that $\frac{1}{L}F + (1 - \frac{1}{L})I$ is nonexpansive, x_* a fixed point of F , $x_0 \in \mathcal{X}$ such that $F(x_0) \neq x_0$, $\gamma > 0$ and for $t \geq 0$,*

$$x_{t+1} = x_t + \frac{\gamma}{\sqrt{\sum_{t=0}^T \|F(x_t) - x_t\|_2^2}} (F(x_t) - x_t).$$

Let $T \geq 0$.

(i) Then,

$$\min_{0 \leq t \leq T} \|F(x_t) - x_t\|_2 \leq \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right) \frac{2L}{\sqrt{T}}.$$

(ii) If $R \geq \max_{0 \leq t \leq T} \|x_t - x_*\|_2$, then $\gamma = R/\sqrt{2}$ yields

$$\min_{0 \leq t \leq T} \|F(x_t) - x_t\|_2 \leq \frac{2\sqrt{2}RL}{\sqrt{T}}.$$

Proof. Combining Lemma 8.4.8 with the regret bound for AdaGrad-Norm (Proposition 7.2.3) gives

$$\sum_{t=0}^T \|F(x_t) - x_t\|_2^2 \leq 2L \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right) \sqrt{\sum_{t=0}^T \|F(x_t) - x_t\|_2^2}.$$

Dividing by $\sqrt{\sum_{t=0}^T \|F(x_t) - x_t\|_2^2}$ gives

$$\sqrt{\sum_{t=0}^T \|F(x_t) - x_t\|_2^2} \leq 2L \left(\frac{\max_{0 \leq t \leq T} \|x_t - x_*\|_2^2}{2\gamma} + \gamma \right).$$

The result follows. \square

Chapter 9

Regret learning in games

For $d \geq 1$, recall that the simplex in \mathbb{R}^d is defined as

$$\Delta_d = \left\{ x \in \mathbb{R}_+^d, \sum_{i=1}^d x_i = 1 \right\},$$

which represents the set of probability distributions over the set $\{1, \dots, d\}$.

9.1 Normal-form games

We quickly recall basic definitions about finite normal-form games.

Definition 9.1.1. A *finite (normal-form) game* is given by

- a finite number of players $N \geq 1$,
- and for each player $1 \leq k \leq N$,
 - a set of $\mathcal{S}^{(k)}$ of *pure strategies*,
 - a payoff function $g^{(k)} : \prod_{k'=1}^N \mathcal{S}^{(k')} \rightarrow \mathbb{R}$

The interpretation is the following. Each player $1 \leq k \leq N$ chooses a strategy $s^{(k)} \in \mathcal{S}^{(k)}$, independently of the other players, and gets payoff $g^{(k)}(s^{(1)}, \dots, s^{(N)})$. Each player aims at maximizing his payoff.

Let such a game be given.

Definition 9.1.2. A N -tuple $(s^{(1)}, \dots, s^{(N)}) \in \mathcal{S}^{(1)} \times \dots \times \mathcal{S}^{(N)}$ is a *Nash equilibrium in pure strategies* if for all $1 \leq k \leq N$,

$$s^{(k)} \in \underset{\tilde{s}^{(k)} \in \mathcal{S}^{(k)}}{\text{Arg max}} g^{(k)}(s^{(1)}, \dots, s^{(k-1)}, \tilde{s}^{(k)}, s^{(k+1)}, \dots, s^{(N)}).$$

Example 9.1.3 (Penalty game). A goalkeeper (Player 1) plays against the shooter (Player 2). Each player chooses left or right. The goalkeeper (resp.

the shooter) wins (resp. loses) if their choices match. $K = 2$, $\mathcal{S}^{(1)} = \mathcal{S}^{(2)} = \{L, R\}$, and

$$g^{(1)}(s^{(1)}, s^{(2)}) = -g^{(2)}(s^{(1)}, s^{(2)}) = \begin{cases} 1 & \text{if } s^{(1)} = s^{(2)} \\ -1 & \text{otherwise.} \end{cases}$$

The game has no Nash equilibrium in pure strategies.

Definition 9.1.4. Let $1 \leq k \leq N$. A *mixed strategy* for Player k , is a probability distribution $a^{(k)} \in \Delta(\mathcal{S}^{(k)})$ over $\mathcal{S}^{(k)}$, where $\Delta(\mathcal{S}^{(k)})$ is the unit simplex in $\mathbb{R}^{\mathcal{S}^{(k)}}$.

Definition 9.1.5. The *mixed extension* of the game from Definition 9.1.1 is the game with sets of strategies $\Delta(\mathcal{S}^{(k)})$ ($1 \leq k \leq N$) and payoff functions given for $1 \leq k \leq N$ and $a^{(1)} \in \Delta(\mathcal{S}^{(1)}), \dots, a^{(N)} \in \Delta(\mathcal{S}^{(N)})$ by

$$g^{(k)}(a^{(1)}, \dots, a^{(N)}) = \sum_{\substack{s^{(1)} \in \mathcal{S}^{(1)} \\ \vdots \\ s^{(N)} \in \mathcal{S}^{(N)}}} \left(\prod_{k'=1}^N a_{s^{(k')}}^{(k')} \right) g^{(k)}(s^{(1)}, \dots, s^{(N)}).$$

Interpretation of the mixed extension is as follows. Each player $1 \leq k \leq N$ independently chooses a mixed action $a^{(k)} \in \Delta(\mathcal{S}^{(k)})$ and draws a pure action $s^{(k)} \sim a^{(k)}$. We then consider the expectation of each payoff $g^{(k)}(s^{(1)}, \dots, s^{(N)})$ ($1 \leq k \leq N$).

Definition 9.1.6. A N -tuple $(a^{(1)}, \dots, a^{(N)}) \in \Delta(\mathcal{S}^{(1)}) \times \dots \times \Delta(\mathcal{S}^{(N)})$ is a *Nash equilibrium in mixed strategies* if for all $1 \leq k \leq N$,

$$a^{(k)} \in \operatorname{Arg\,max}_{\tilde{a}^{(k)} \in \Delta(\mathcal{S}^{(k)})} g_k(a^{(1)}, \dots, a^{(k-1)}, \tilde{a}^{(k)}, a^{(k+1)}, \dots, a^{(N)}).$$

Theorem 9.1.7 (Nash, 1951). *A finite game admits a Nash equilibrium in mixed strategies.*

Example 9.1.8 (Penalty game). In the penalty game, each player choosing left or right with probability 1/2 is a Nash equilibrium.

9.2 Two-player zero-sum games

Let $m, n \geq 1$ be integers. We focus on zero-sum two-player games which can be represented by a matrix $A \in \mathbb{R}^{m \times n}$. Denote $\mathcal{S}^{(1)} = \{1, \dots, m\}$ (resp. $\mathcal{S}^{(2)} = \{1, \dots, n\}$) the set of *pure strategies* of Player 1 (resp. Player 2). When Player 1 and Player 2 choose strategies $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$ respectively, Player 1 (resp. Player 2) obtains payoff A_{ij} (resp. $-A_{ij}$). For mixed strategies $a \in \Delta_m$ and $b \in \Delta_n$, if $i \sim a$ and $j \sim b$,

$$\mathbb{E}[A_{ij}] = \sum_{i=1}^m \sum_{j=1}^n a_i b_j A_{ij} = \langle a, Ab \rangle.$$

Remark 9.2.1. $(a_*, b_*) \in \Delta_m \times \Delta_n$ is a Nash equilibrium in mixed strategies if, and only if,

$$a_* \in \operatorname{Arg} \max_{a \in \Delta_m} \langle a, Ab_* \rangle \quad \text{and} \quad b_* \in \operatorname{Arg} \min_{b \in \Delta_n} \langle a_*, Ab \rangle.$$

Definition 9.2.2. The *duality gap* $\delta_A : \Delta_m \times \Delta_n \rightarrow \mathbb{R}_+$ is defined as

$$\delta_A(a, b) = \max_{a' \in \Delta_m} \langle a', Ab \rangle - \min_{b' \in \Delta_n} \langle a, Ab' \rangle, \quad a \in \Delta_m, b \in \Delta_n.$$

Proposition 9.2.3. $(a_*, b_*) \in \Delta_m \times \Delta_n$ is a Nash equilibrium of the two-player zero-sum game associated with A if, and only if, $\delta_A(a_*, b_*) = 0$.

Proof. It always holds that

$$\max_{a \in \Delta_m} \langle a, Ab_* \rangle \geq \langle a_*, Ab_* \rangle \geq \min_{b \in \Delta_n} \langle a_*, Ab \rangle.$$

Therefore, $\delta_A(a_*, b_*) = 0$ if, and only, if the above inequalities are equalities. The above first inequality being an equality is equivalent to

$$a_* \in \operatorname{Arg} \max_{a \in \Delta_m} \langle a, Ab_* \rangle,$$

and similarly the second inequality being an equality is equivalent to

$$b_* \in \operatorname{Arg} \min_{b \in \Delta_n} \langle a_*, Ab \rangle.$$

Hence the result. □

Remark 9.2.4. The duality gap is easy to compute, as it rewrites

$$\delta_A(a, b) = \max_{1 \leq i \leq m} (Ab)_i - \min_{1 \leq j \leq n} (A^\top a)_j, \quad a \in \Delta_m, b \in \Delta_n.$$

It is therefore the quantity of choice to measure how far a couple (a, b) is from being a Nash equilibrium.

Theorem 9.2.5 (Von Neumann's minimax theorem). *There exists a Nash equilibrium $(a_*, b_*) \in \Delta_m \times \Delta_n$ of the two-player zero-sum game associated with A and*

$$\langle a_*, Ab_* \rangle = \max_{a \in \Delta_m} \min_{b \in \Delta_n} \langle a, Ab \rangle = \min_{b \in \Delta_n} \max_{a \in \Delta_m} \langle a, Ab \rangle.$$

There are several proofs of von Neumann's theorem. In the next section, we give one based on regret minimization.

9.3 Regret learning

The following statement demonstrates that the duality gap can be minimized if the game is played repeatedly, with each player minimizing its regret.

Lemma 9.3.1. *Let $T \geq 0$, $a_0, \dots, a_T \in \Delta_m$, $b_0, \dots, b_T \in \Delta_n$, $\bar{a}_T = \frac{1}{T+1} \sum_{t=0}^T a_t$ and $\bar{b}_T = \frac{1}{T+1} \sum_{t=0}^T b_t$. Then,*

$$\delta_T(\bar{a}_T, \bar{b}_T) = \frac{1}{T+1} \left(\max_{a \in \Delta_m} \sum_{t=0}^T \langle Ab_t, a - a_t \rangle + \max_{b \in \Delta_n} \sum_{t=0}^T \langle -A^\top a_t, b - b_t \rangle \right).$$

Proof.

$$\begin{aligned} \delta_T(\bar{a}_T, \bar{b}_T) &= \max_{a \in \Delta_m} \left\langle a, A \left(\frac{1}{T+1} \sum_{t=0}^T b_t \right) \right\rangle - \min_{b \in \Delta_n} \left\langle \frac{1}{T+1} \sum_{t=0}^T a_t, Ab \right\rangle \\ &= \frac{1}{T+1} \left(\max_{a \in \Delta_m} \sum_{t=0}^T \langle a, Ab_t \rangle - \max_{b \in \Delta_n} \sum_{t=0}^T \langle -A^\top a_t, b \rangle \right) \\ &= \frac{1}{T+1} \left(\max_{a \in \Delta_m} \sum_{t=0}^T \langle a, Ab_t \rangle - \sum_{t=0}^T \langle a_t, Ab_t \rangle + \sum_{t=0}^T \langle A^\top a_t, b_t \rangle \right. \\ &\quad \left. + \max_{b \in \Delta_n} \sum_{t=0}^T \langle -A^\top a_t, b \rangle \right), \end{aligned}$$

hence the result. \square

The following proposition uses the exponential weights algorithm to minimize each of the two regrets appearing in the above lemma. The guarantee on the duality gap is then an immediate adaptation from the regret bound for the exponential weights algorithm (Proposition 3.4.4).

Proposition 9.3.2 (Exponential weights for two-player zero-sum games).

For $t \geq 0$, let

$$y_t = \eta_t \sum_{s=0}^{t-1} Ab_s, \quad z_t = -\eta'_t \sum_{s=0}^{t-1} A^\top a_s,$$

where

$$\eta_t = \frac{\sqrt{2 \log m}}{\|A\|_\infty \sqrt{t+1}}, \quad \eta'_t = \frac{\sqrt{2 \log n}}{\|A\|_\infty \sqrt{t+1}},$$

and

$$a_t = \left(\frac{\exp(y_{t,i})}{\sum_{i'=1}^m \exp(y_{t,i'})} \right)_{1 \leq i \leq m}, \quad b_t = \left(\frac{\exp(z_{t,j})}{\sum_{j'=1}^n \exp(z_{t,j'})} \right)_{1 \leq j \leq n},$$

For $T \geq 0$, let $\bar{a}_T = \frac{1}{T+1} \sum_{t=0}^T a_t$ and $\bar{b}_T = \frac{1}{T+1} \sum_{t=0}^T b_t$. Then,

$$\delta_A(\bar{a}_T, \bar{b}_T) \leq \|A\|_\infty \left(\sqrt{\log m} + \sqrt{\log n} \right) \sqrt{\frac{2}{T+1}}.$$

Proof. For all $t \geq 0$, $\|Ab_t\|_\infty \leq \|A\|_\infty \|b_t\|_1 = \|A\|_\infty$ because $b_t \in \Delta_n$. Similarly, $\|A^\top a_t\|_\infty \leq \|A\|_\infty$. Therefore, similarly to Proposition 3.4.4, we obtain regret bounds

$$\begin{aligned} \max_{a \in \Delta_m} \sum_{t=0}^T \langle Ab_t, a - a_t \rangle &\leq \|A\|_\infty \sqrt{2(\log m)(T+1)} \\ \max_{b \in \Delta_n} \sum_{t=0}^T \langle -A^\top a_t, b - b_t \rangle &\leq \|A\|_\infty \sqrt{2(\log n)(T+1)}. \end{aligned}$$

Summing and applying Lemma 9.3.1 gives the result. \square

We can use the above result to now prove von Neumann's minmax theorem.

Proof of Theorem 9.2.5. Let us first prove that δ_A is continuous on $\Delta_m \times \Delta_n$.

$$(a, b) \mapsto \max_{a' \in \Delta_m} \langle a', Ab \rangle = \max_{a' \in \Delta_m} \langle A^\top a', b \rangle,$$

is defined on $\mathbb{R}^m \times \mathbb{R}^n$ and is convex as the point-wise maximum of linear functions. It is therefore continuous. Similarly, $(a, b) \mapsto \max_{b' \in \Delta_n} \langle a, Ab' \rangle$ is continuous. Hence, δ_A is continuous.

For all $T \geq 0$, let $\bar{a}_T \in \Delta_m$ and $\bar{b}_T \in \Delta_n$ from Proposition 9.3.2. Because they belong to simplexes which are compact sets, there exists subsequences so that they converge to $a_* \in \Delta_m$ and $b_* \in \Delta_n$ respectively. Because the bound from Proposition 9.3.2 is vanishing as $T \rightarrow \infty$, by continuity of δ_A , $\delta_A(a_*, b_*) = 0$, in other words, (a_*, b_*) is a Nash equilibrium by Proposition 9.2.3.

Moreover, for all $(a, b) \in \Delta_m \times \Delta_n$, in addition to

$$\max_{a' \in \Delta_m} \langle a', Ab \rangle \geq \langle a, Ab \rangle \geq \min_{b' \in \Delta_n} \langle a, Ab' \rangle,$$

it is elementary to prove that

$$\max_{a'} \langle a', Ab \rangle \geq \min_{b' \in \Delta_n} \max_{a' \in \Delta_m} \langle a', Ab' \rangle \geq \max_{a' \in \Delta_m} \min_{b' \in \Delta_n} \langle a', Ab' \rangle \geq \min_{b' \in \Delta_n} \langle a, Ab' \rangle.$$

For $a = a_*$ and $b = b_*$ in particular, because $\delta_A(a_*, b_*) = 0$, the left-most and right-most quantities of both above displays are equal. Therefore, all inequalities are equalities and

$$\langle a_*, Ab_* \rangle = \min_{b' \in \Delta_n} \max_{a' \in \Delta_m} \langle a', Ab' \rangle = \max_{a' \in \Delta_m} \min_{b' \in \Delta_n} \langle a', Ab' \rangle.$$

\square

Proposition 9.3.3 (RM for two-player zero-sum games). *Let $a_0 \in \Delta_m$ and $b_0 \in \Delta_n$. For $t \geq 0$, let*

$$a_t = \begin{cases} \frac{x_{t,+}}{\|x_{t,+}\|_1} & \text{if } x_{t,+} := \left(\sum_{s=0}^{t-1} (Ab_s - \langle a_s, Ab_s \rangle \mathbb{1}) \right)_+ \neq 0 \\ a_0 & \text{otherwise,} \end{cases}$$

$$b_t = \begin{cases} \frac{w_{t,+}}{\|w_{t,+}\|_1} & \text{if } w_{t,+} := \left(\sum_{s=0}^{t-1} (\langle a_s, Ab_s \rangle \mathbb{1} - A^\top a_s) \right)_+ \neq 0 \\ b_0 & \text{otherwise.} \end{cases}$$

For all $T \geq 0$, let $\bar{a}_T = \frac{1}{T+1} \sum_{t=0}^T a_t$ and $\bar{b}_T = \frac{1}{T+1} \sum_{t=0}^T b_t$. Then,

$$\delta_A(\bar{a}_T, \bar{b}_T) \leq (A_{\max} - A_{\min}) \frac{\sqrt{m} + \sqrt{n}}{\sqrt{T+1}}.$$

Proof. The general regret bound for RM from Proposition 5.5.3 gives

$$\max_{a \in \Delta_m} \sum_{t=0}^T \langle Ab_t, a - a_t \rangle \leq \sqrt{\sum_{t=0}^T \|Ab_t - \langle a_t, Ab_t \rangle \mathbb{1}\|_2^2}$$

$$\max_{b \in \Delta_n} \sum_{t=0}^T \langle -A^\top a_t, b - b_t \rangle \leq \sqrt{\sum_{t=0}^T \|\langle a_t, Ab_t \rangle \mathbb{1} - A^\top a_t\|_2^2}$$

For all $t \geq 0$, because $a_t \in \Delta_m$ and $b_t \in \Delta_n$,

$$\begin{aligned} \|Ab_t - \langle a_t, Ab_t \rangle \mathbb{1}\|_2^2 &= \sum_{i=1}^m ((Ab_t)_i - \langle a_t, Ab_t \rangle)^2 \\ &\leq \sum_{i=1}^m (A_{\max} - A_{\min})^2 \\ &= m (A_{\max} - A_{\min})^2, \end{aligned}$$

and similarly

$$\|\langle a_t, Ab_t \rangle \mathbb{1} - A^\top a_t\|_2^2 \leq n (A_{\max} - A_{\min})^2.$$

The result then follows by applying Lemma 9.3.1. \square

Proposition 9.3.4 (RM+ for two-player zero-sum games). *Let $a_0 \in \Delta_m$, $b_0 \in \Delta_n$, $x_0 = 0$, $w_0 = 0$. For $t \geq 0$, let*

$$a_t = \begin{cases} \frac{x_t}{\|x_t\|_1} & \text{if } x_t \neq 0 \\ a_0 & \text{otherwise,} \end{cases} \quad b_t = \begin{cases} \frac{w_t}{\|w_t\|_1} & \text{if } w_t \neq 0 \\ b_0 & \text{otherwise.} \end{cases}$$

$$x_{t+1} = (x_t + Ab_t - \langle a_t, Ab_t \rangle \mathbb{1})_+ \quad w_{t+1} = (w_t + \langle a_t, Ab_t \rangle \mathbb{1} - A^\top a_t)_+.$$

Then, the same guarantee as for RM (Proposition 9.3.3) holds.

Proof. Identical to Proposition 9.3.3. \square

9.4 Optimistic regret learning

Section 8.3 provided an approach for solving a class of problems containing two-player zero-sum games with fast convergence rate $1/T$. We here provide an alternative approach to obtain such a rate.

Informally, an optimistic variant of a regret minimization algorithm uses the last observed vector twice, the second time as a guess for the next vector. For instance, an optimistic exponential weights algorithm against $(u_t)_{t \geq 0}$ would write

$$x_t = \left(\frac{\exp\left(\eta \sum_{s=0}^{t-1} u_{s,i} + u_{t-1,i}\right)}{\sum_{i'=1}^d \exp\left(\eta \sum_{s=0}^{t-1} u_{s,i'} + u_{t-1,i'}\right)} \right)_{1 \leq i \leq d} \quad t \geq 0.$$

Lemma 9.4.1 (Optimistic UMD). *Let $\|\cdot\|$ be a norm on \mathbb{R}^d , $K > 0$, \mathcal{X} be a nonempty closed convex set, h a regularizer on \mathcal{X} which is K -strongly convex for $\|\cdot\|$, $(u_t)_{t \geq 0}$ a sequence in \mathbb{R}^d , and $((x_t, y_t))_{t \geq 0}$ be a sequence of strict UMD iterates associated with regularizer h and dual iterates $(2u_t - u_{t-1})_{t \geq 0}$ (with convention $u_{-1} = 0$). Then for all $T \geq 0$, $\alpha > 0$, and $x \in \text{dom } h$,*

$$\begin{aligned} \sum_{t=0}^T \langle u_t, x - x_t \rangle &\leq D_h(x, x_0; y_0) + \frac{\alpha}{2} \|x - x_T\|^2 + \frac{1}{2\alpha} \|u_0\|_*^2 \\ &\quad + \frac{1}{2\alpha} \sum_{t=0}^{T-1} \|u_{t+1} - u_t\|_*^2 + \left(\frac{\alpha}{2} - \frac{K}{2}\right) \sum_{t=0}^{T-1} \|x_{t+1} - x_t\|^2. \end{aligned}$$

Proof.

$$\begin{aligned} \sum_{t=0}^T \langle u_t, x - x_t \rangle &= \langle u_0, x - x_0 \rangle + \sum_{t=0}^{T-1} \langle u_{t+1} - 2u_t + u_{t-1}, x - x_{t+1} \rangle \\ &\quad + \sum_{t=0}^{T-1} \langle 2u_t - u_{t-1}, x - x_{t-1} \rangle. \end{aligned} \tag{9.1}$$

The above last sum can be bounded from above using Lemma 2.4.1 and the strong convexity of h :

$$\begin{aligned} \sum_{t=0}^{T-1} \langle 2u_t - u_{t-1}, x - x_{t+1} \rangle &\leq D_h(x, x_0; y_0) - \sum_{t=0}^{T-1} D_h(x_{t+1}, x_t; y_t) \\ &\leq D_h(x, x_0; y_0) - \frac{K}{2} \sum_{t=0}^{T-1} \|x_{t+1} - x_t\|^2. \end{aligned} \tag{9.2}$$

The other sum is bounded as follows.

$$\begin{aligned}
\sum_{t=0}^{T-1} \langle u_{t+1} - 2u_t + u_{t-1}, x - x_{t+1} \rangle &= \sum_{t=0}^{T-1} \langle u_{t+1} - u_t, x - x_{t+1} \rangle \\
&\quad - \sum_{t=0}^{T-1} \langle u_t - u_{t-1}, x - x_{t+1} \rangle \\
&= \sum_{t=0}^{T-1} \langle u_{t+1} - u_t, x - x_{t+1} \rangle - \sum_{t=0}^{T-2} \langle u_{t+1} - u_t, x - x_{t+2} \rangle - \langle u_0, x - x_1 \rangle \\
&= \sum_{t=0}^{T-2} \langle u_{t+1} - u_t, x_{t+2} - x_{t+1} \rangle + \langle u_T - u_{T-1}, x - x_T \rangle - \langle u_0, x - x_1 \rangle \\
&\leq \sum_{t=0}^{T-2} \left(\frac{\alpha}{2} \|x_{t+2} - x_{t+1}\|^2 + \frac{1}{2\alpha} \|u_{t+1} - u_t\|_*^2 \right) \\
&\quad + \frac{1}{2\alpha} \|u_T - u_{T-1}\|_*^2 + \frac{\alpha}{2} \|x - x_T\|^2 - \langle u_0, x - x_1 \rangle.
\end{aligned} \tag{9.3}$$

Combining (9.1), (9.2) and (9.3) gives

$$\begin{aligned}
\sum_{t=0}^T \langle u_t, x - x_t \rangle &\leq D_h(x, x_0; y_0) + \langle u_0, x_1 - x_0 \rangle + \frac{\alpha}{2} \sum_{t=0}^{T-2} \|x_{t+2} - x_{t+1}\|^2 \\
&\quad - \frac{K}{2} \sum_{t=0}^{T-1} \|x_{t+1} - x_t\|^2 + \frac{\alpha}{2} \|x - x_T\|^2 \\
&\quad + \frac{1}{2\alpha} \sum_{t=0}^{T-1} \|u_{t+1} - u_t\|_*^2.
\end{aligned}$$

Regarding the second term in the above upper bound, we write

$$\langle u_0, x_1 - x_0 \rangle \leq \frac{1}{2\alpha} \|u_0\|_*^2 + \frac{\alpha}{2} \|x_1 - x_0\|^2.$$

The result follows. \square

Proposition 9.4.2 (Optimistic exponential weights for two-player zero-sum games). *Let $\eta > 0$ and for $t \geq 0$, let*

$$y_t = \eta \left(\sum_{s=0}^{t-1} Ab_s + Ab_{t-1} \right), \quad z_t = -\eta \left(\sum_{s=0}^{t-1} A^\top a_s + A^\top a_{t-1} \right),$$

and

$$a_t = \left(\frac{\exp(y_{t,i})}{\sum_{i'=1}^m \exp(y_{t,i'})} \right)_{1 \leq i \leq m}, \quad b_t = \left(\frac{\exp(z_{t,j})}{\sum_{j'=1}^m \exp(z_{t,j'})} \right)_{1 \leq j \leq n}.$$

- (i) $(a_t)_{t \geq 0}$ (resp. $(b_t)_{t \geq 0}$) corresponds to the exponential weights algorithm associated with dual increments $(\eta(2Ab_t - Ab_{t-1}))_{t \geq 0}$ (resp. $(\eta(2A^\top a_t - A^\top a_{t-1}))_{t \geq 0}$) (with convention $b_{-1} = 0$ and $a_{-1} = 0$).
- (ii) For $T \geq 0$, let $\bar{a}_T = \frac{1}{T+1} \sum_{t=0}^T a_t$ and $\bar{b}_T = \frac{1}{T+1} \sum_{t=0}^T b_t$. Then, if $\eta = 1/(2\|A\|_\infty)$,

$$\delta_A(\bar{a}_T, \bar{b}_T) \leq \frac{2\|A\|_\infty (\log m + \log n + 2)}{T+1}.$$

Proof. It is easy to verify that for all $t \geq 0$, $y_t = \eta \sum_{s=0}^{t-1} (2Ab_t - Ab_{t-1})$. Similarly $z_t = -\eta \sum_{s=0}^{t-1} (2A^\top a_t - A^\top a_{t-1})$. Hence (i) holds.

Because $y_0 = 0$, if h is the entropic regularizer on Δ_m ,

$$D_h(a, a_0; y_0) = h(a) - h(a_0) \leq \log m,$$

by Proposition 3.4.3. Similarly, if \tilde{h} is the entropic regularizer on Δ_n , because $z_0 = 0$,

$$D_{\tilde{h}}(b, b_0; z_0) \leq \log n.$$

Then, because the entropic regularizer is 1-strongly convex for $\|\cdot\|_1$ by Proposition 2.2.6, applying Lemma 9.4.1 gives regret bounds

$$\begin{aligned} \eta \sum_{t=0}^T \langle Ab_t, a - a_t \rangle &\leq \log m + \frac{1}{4} \|a - a_T\|_1^2 + \|2\eta Ab_0\|_\infty^2 \\ &\quad + \eta^2 \sum_{t=0}^{T-1} \|Ab_{t+1} - Ab_t\|_\infty^2 - \frac{1}{4} \sum_{t=0}^{T-1} \|a_{t+1} - a_t\|_1^2, \\ \eta \sum_{t=0}^T \langle -A^\top a_t, b - b_t \rangle &\leq \log n + \frac{1}{4} \|b - b_T\|_1^2 + \|2\eta A^\top a_0\|_\infty^2 \\ &\quad + \eta^2 \sum_{t=0}^{T-1} \|A^\top a_{t+1} - A^\top a_t\|_\infty^2 - \frac{1}{4} \sum_{t=0}^{T-1} \|b_{t+1} - b_t\|_1^2. \end{aligned}$$

Note that for all $t \geq 0$,

$$\begin{aligned} \|Ab_{t+1} - Ab_t\|_\infty^2 &\leq \|A\|_\infty^2 \|b_{t+1} - b_t\|_1^2, \\ \|A^\top a_{t+1} - A^\top a_t\|_\infty^2 &\leq \|A\|_\infty^2 \|a_{t+1} - a_t\|_1^2. \end{aligned}$$

Besides,

$$\|2Ab_0\|_\infty^2 \leq 4\|A\|_\infty^2 \|b_0\|_1^2 = 4\|A\|_\infty^2,$$

similarly $\|2A^\top a_0\|_\infty^2 \leq 4\|A\|_\infty^2$, and

$$\|a - a_T\|_1^2 \leq (\|a\|_1 + \|a_T\|_1)^2 = 4,$$

and similarly $\|b - b_T\|_1^2 \leq 4$.

Dividing by η , summing, and applying Lemma 9.3.1 with $\alpha = 1/2$ gives

$$(T+1) \cdot \delta_A(\bar{a}_T, \bar{b}_T) \leq \frac{\log m + \log n + 2}{\eta} + 8\eta \|A\|_\infty^2 \\ + \left(\eta \|A\|_\infty^2 - \frac{1}{4\eta} \right) \sum_{t=0}^{T-1} \left(\|a_{t+1} - a_t\|_1^2 + \|b_{t+1} - b_t\|_1^2 \right).$$

Because $\eta = 1/(2\|A\|_\infty)$, the above last term is zero and the result follows. \square

Bibliography

- [Bub11] Sébastien Bubeck. *Introduction to Online Optimization: Lecture Notes*. Princeton University, 2011.
- [CBL06] Nicolò Cesa-Bianchi and Gábor Lugosi. *Prediction, Learning, and Games*. Cambridge University Press, 2006.
- [Haz16] Elad Hazan. Introduction to online convex optimization. *Foundations and Trends® in Optimization*, 2(3-4):157–325, 2016.
- [JKM23] Anatoli Juditsky, Joon Kwon, and Éric Moulines. Unifying mirror descent and dual averaging. *Mathematical Programming*, (199):793–830, 2023.
- [McM17] H. Brendan McMahan. A survey of algorithms and analysis for adaptive online learning. *The Journal of Machine Learning Research*, 18(1):3117–3166, 2017.
- [Ora23] Francesco Orabona. A modern introduction to online learning. *arXiv:1912.13213*, 2023.
- [Roc70] R. Tyrrell Rockafellar. *Convex Analysis*. Princeton University Press, 1970.
- [SS11] Shai Shalev-Shwartz. Online learning and online convex optimization. *Foundations and Trends in Machine Learning*, 4(2):107–194, 2011.